

Submodular Functions, Optimization, and Applications to Machine Learning

— Spring Quarter, Lecture 15 —

http://www.ee.washington.edu/people/faculty/bilmes/classes/ee563_spring_2018/

Prof. Jeff Bilmes

University of Washington, Seattle
Department of Electrical Engineering
<http://melodi.ee.washington.edu/~bilmes>

May 16th, 2018



$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$$

$$-f(A) + 2f(C) + f(B) \quad -f(A) + f(C) + f(B) \quad -f(A \cap B)$$



Cumulative Outstanding Reading

- Read chapter 1 from Fujishige's book.
- Read chapter 2 from Fujishige's book.
- Read chapter 3 from Fujishige's book.
- Read chapter 4 from Fujishige's book.

Announcements, Assignments, and Reminders

- Next homework will be posted soon.
- As always, if you have any questions about anything, please ask them via our discussion board (https://canvas.uw.edu/courses/1216339/discussion_topics). Can meet at odd hours via zoom (send message on canvas to schedule time to chat).

Class Road Map - EE563

- L1(3/26): Motivation, Applications, & Basic Definitions,
- L2(3/28): Machine Learning Apps (diversity, complexity, parameter, learning target, surrogate).
- L3(4/2): Info theory exs, more apps, definitions, graph/combinatorial examples
- L4(4/4): Graph and Combinatorial Examples, Matrix Rank, Examples and Properties, visualizations
- L5(4/9): More Examples/Properties/ Other Submodular Defs., Independence,
- L6(4/11): Matroids, Matroid Examples, Matroid Rank, Partition/Laminar Matroids
- L7(4/16): Laminar Matroids, System of Distinct Reps, Transversals, Transversal Matroid, Matroid Representation, Dual Matroids
- L8(4/18): Dual Matroids, Other Matroid Properties, Combinatorial Geometries, Matroids and Greedy.
- L9(4/23): Polyhedra, Matroid Polytopes, Matroids \rightarrow Polymatroids
- L10(4/29): Matroids \rightarrow Polymatroids, Polymatroids, Polymatroids and Greedy,
- L11(4/30): Polymatroids, Polymatroids and Greedy
- L12(5/2): Polymatroids and Greedy, Extreme Points, Cardinality Constrained Maximization
- L13(5/7): Constrained Submodular Maximization
- L14(5/9): Submodular Max w. Other Constraints, Cont. Extensions, Lovasz Extension
- L15(5/14): Cont. Extensions, Lovasz Extension, Choquet Integration, Properties
- L16(5/16):
- L17(5/21):
- L18(5/23):
- L-(5/28): Memorial Day (holiday)
- L19(5/30):
- L21(6/4): Final Presentations maximization.

Last day of instruction, June 1st. Finals Week: June 2-8, 2018.

Continuous Extensions of Discrete Set Functions

- Any function $f : 2^V \rightarrow \mathbb{R}$ (equivalently $f : \{0, 1\}^V \rightarrow \mathbb{R}$) can be extended to a continuous function in the sense $\tilde{f} : [0, 1]^V \rightarrow \mathbb{R}$.
- This may be tight (i.e., $\tilde{f}(\mathbf{1}_A) = f(A)$ for all A). I.e., the extension \tilde{f} coincides with f at the hypercube vertices.
- In fact, any such discrete function defined on the vertices of the n -D hypercube $\{0, 1\}^n$ has a variety of both convex and concave extensions tight at the vertices (Crama & Hammer'11). Example $n = 1$,

Concave Extensions


$$\tilde{f} : [0, 1] \rightarrow \mathbb{R}$$

Discrete Function

$$f : \{0, 1\}^V \rightarrow \mathbb{R}$$

Convex Extensions

$$\tilde{f} : [0, 1] \rightarrow \mathbb{R}$$



- Since there are an exponential number of vertices $\{0, 1\}^n$, important questions regarding such extensions is:
 - When are they computationally feasible to obtain or estimate?
 - When do they have nice mathematical properties?
 - When are they useful for something practical?

Def: Convex Envelope of a function


- Given any function $h : \mathbb{R}^n \rightarrow \mathbb{R}$, define new function $\check{h} : \mathbb{R}^n \rightarrow \mathbb{R}$ via:

$$\check{h}(x) = \sup \{g(x) : g \text{ is convex} \ \& \ g(y) \leq h(y), \forall y \in \mathbb{R}^n\} \quad (15.6)$$

$\forall y \in D_h \subseteq \mathbb{R}^n$

- I.e., (1) $\check{h}(x)$ is convex, (2) $\check{h}(x) \leq h(x)$, $\forall x$, and (3) if $g(x)$ is any convex function having the property that $g(x) \leq h(x)$, $\forall x$, then $g(x) \leq \check{h}(x)$.
- Alternatively,

$$\check{h}(x) = \inf \{t : (x, t) \in \text{convexhull}(\text{epigraph}(h))\} \quad (15.7)$$




Convex Closure of Discrete Set Functions

- Given set function $f : 2^V \rightarrow \mathbb{R}$, an arbitrary (i.e., not necessarily submodular nor supermodular) set function, define a function $\check{f} : [0, 1]^V \rightarrow \mathbb{R}$, as

$$\check{f}(x) = \min_{p \in \Delta^n(x)} \sum_{S \subseteq V} p_S f(S) \quad (15.1)$$

where $\Delta^n(x) =$

$$\left\{ p \in \mathbb{R}^{2^n} : \sum_{S \subseteq V} p_S = 1, p_S \geq 0 \forall S \subseteq V, \& \sum_{S \subseteq V} p_S \mathbf{1}_S = x \right\}$$



$$\forall p \in \Delta^2(x)$$

$$x = p_\emptyset \cdot \mathbf{1}_\emptyset + p_{v_1} \cdot \mathbf{1}_{\{v_1\}} + p_{v_2} \cdot \mathbf{1}_{\{v_2\}} + p_{v_1, v_2} \cdot \mathbf{1}_{\{v_1, v_2\}}$$

Convex Closure of Discrete Set Functions

- Given set function $f : 2^V \rightarrow \mathbb{R}$, an arbitrary (i.e., not necessarily submodular nor supermodular) set function, define a function $\check{f} : [0, 1]^V \rightarrow \mathbb{R}$, as

$$\check{f}(x) = \min_{p \in \Delta^n(x)} \sum_{S \subseteq V} p_S f(S) \quad (15.1)$$

where $\Delta^n(x) =$

$$\left\{ p \in \mathbb{R}^{2^n} : \sum_{S \subseteq V} p_S = 1, p_S \geq 0 \forall S \subseteq V, \& \sum_{S \subseteq V} p_S \mathbf{1}_S = x \right\}$$

- Hence, $\Delta^n(x)$ is the set of all probability distributions over the 2^n vertices of the hypercube, and where the expected value of the characteristic vectors of those points is equal to x , i.e., for any $p \in \Delta^n(x)$, $E_{S \sim p}(\mathbf{1}_S) = \sum_{S \subseteq V} p_S \mathbf{1}_S = x$.

Convex Closure of Discrete Set Functions

- Given set function $f : 2^V \rightarrow \mathbb{R}$, an arbitrary (i.e., not necessarily submodular nor supermodular) set function, define a function $\check{f} : [0, 1]^V \rightarrow \mathbb{R}$, as

$$\check{f}(x) = \min_{p \in \Delta^n(x)} \sum_{S \subseteq V} p_S f(S) \quad (15.1)$$

where $\Delta^n(x) =$

$$\left\{ p \in \mathbb{R}^{2^n} : \sum_{S \subseteq V} p_S = 1, p_S \geq 0 \forall S \subseteq V, \& \sum_{S \subseteq V} p_S \mathbf{1}_S = x \right\}$$

- Hence, $\Delta^n(x)$ is the set of all probability distributions over the 2^n vertices of the hypercube, and where the expected value of the characteristic vectors of those points is equal to x , i.e., for any $p \in \Delta^n(x)$, $E_{S \sim p}(\mathbf{1}_S) = \sum_{S \subseteq V} p_S \mathbf{1}_S = x$.
- Hence, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$

Convex Closure of Discrete Set Functions

- Given set function $f : 2^V \rightarrow \mathbb{R}$, an arbitrary (i.e., not necessarily submodular nor supermodular) set function, define a function $\check{f} : [0, 1]^V \rightarrow \mathbb{R}$, as

$$\check{f}(x) = \min_{p \in \Delta^n(x)} \sum_{S \subseteq V} p_S f(S) \quad (15.1)$$

where $\Delta^n(x) =$

$$\left\{ p \in \mathbb{R}^{2^n} : \sum_{S \subseteq V} p_S = 1, p_S \geq 0 \forall S \subseteq V, \& \sum_{S \subseteq V} p_S \mathbf{1}_S = x \right\}$$

- Hence, $\Delta^n(x)$ is the set of all probability distributions over the 2^n vertices of the hypercube, and where the expected value of the characteristic vectors of those points is equal to x , i.e., for any $p \in \Delta^n(x)$, $E_{S \sim p}(\mathbf{1}_S) = \sum_{S \subseteq V} p_S \mathbf{1}_S = x$.
- Hence, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$
- Note, this is not (necessarily) the Lovász extension, rather this is a convex extension.

Convex Closure of Discrete Set Functions

- Given, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$, there are several things we'd like to show:

Convex Closure of Discrete Set Functions

- Given, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$, there are several things we'd like to show:
 - That \check{f} is tight (i.e., $\forall S \subseteq V$, we have $\check{f}(\mathbf{1}_S) = f(S)$).

Convex Closure of Discrete Set Functions

- Given, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$, there are several things we'd like to show:
 - That \check{f} is tight (i.e., $\forall S \subseteq V$, we have $\check{f}(\mathbf{1}_S) = f(S)$).
 - That \check{f} is convex (and consequently, that any arbitrary set function has a tight convex extension).

Convex Closure of Discrete Set Functions

- Given, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$, there are several things we'd like to show:
 - That \check{f} is tight (i.e., $\forall S \subseteq V$, we have $\check{f}(\mathbf{1}_S) = f(S)$).
 - That \check{f} is convex (and consequently, that any arbitrary set function has a tight convex extension).
 - That the convex closure \check{f} is the convex envelope of the function defined only on the hypercube vertices, and that takes value $f(S)$ at $\mathbf{1}_S$.

Convex Closure of Discrete Set Functions

- Given, $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$, there are several things we'd like to show:
 - That \check{f} is tight (i.e., $\forall S \subseteq V$, we have $\check{f}(\mathbf{1}_S) = f(S)$).
 - That \check{f} is convex (and consequently, that any arbitrary set function has a tight convex extension).
 - That the convex closure \check{f} is the convex envelope of the function defined only on the hypercube vertices, and that takes value $f(S)$ at $\mathbf{1}_S$.
 - The definition of the Lovász extension of a set function, and that \check{f} is the Lovász extension iff f is submodular.

Tightness of Convex Closure

Lemma 15.3.1

$\forall A \subseteq V$, we have $\check{f}(\mathbf{1}_A) = f(A)$.

Proof.

- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$.

$$p^x \in \operatorname{argmin}_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$$

Tightness of Convex Closure

Lemma 15.3.1

$\forall A \subseteq V$, we have $\check{f}(\mathbf{1}_A) = f(A)$.

Proof.

- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$.
- Take an arbitrary A , so that $\mathbf{1}_A = \sum_{S \subseteq V} p_S^{\mathbf{1}_A} \mathbf{1}_S = \mathbf{1}_A$.


Tightness of Convex Closure

Lemma 15.3.1

$\forall A \subseteq V$, we have $\check{f}(\mathbf{1}_A) = f(A)$.

Proof.

- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$.
- Take an arbitrary A , so that $\mathbf{1}_A = \sum_{S \subseteq V} p_S^{1_A} \mathbf{1}_S = \mathbf{1}_A$.
- Suppose $\exists S'$ with $S' \setminus A \neq \emptyset$ having $p_{S'}^{1_A} > 0$. This would mean, for any $v \in S' \setminus A$, that $(\sum_S p_S^{1_A} \mathbf{1}_S)(v) > 0$, a contradiction.



Tightness of Convex Closure

Lemma 15.3.1

$\forall A \subseteq V$, we have $\check{f}(\mathbf{1}_A) = f(A)$.

Proof.

- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$.
- Take an arbitrary A , so that $\mathbf{1}_A = \sum_{S \subseteq V} p_S^{\mathbf{1}_A} \mathbf{1}_S = \mathbf{1}_A$.
- Suppose $\exists S'$ with $S' \setminus A \neq \emptyset$ having $p_{S'}^{\mathbf{1}_A} > 0$. This would mean, for any $v \in S' \setminus A$, that $\left(\sum_S p_S^{\mathbf{1}_A} \mathbf{1}_S\right)(v) > 0$, a contradiction.
- Suppose $\exists S'$ s.t. $A \setminus S' \neq \emptyset$ with $p_{S'}^{\mathbf{1}_A} > 0$.



Tightness of Convex Closure

Lemma 15.3.1

$\forall A \subseteq V$, we have $\check{f}(\mathbf{1}_A) = f(A)$.

$$E_{S \sim p^A}[f(S)] = 1 \cdot f(A) = \check{f}(\mathbf{1}_A)$$

Proof.

- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$.
- Take an arbitrary A , so that $\mathbf{1}_A = \sum_{S \subseteq V} p_S^{\mathbf{1}_A} \mathbf{1}_S = \mathbf{1}_A$.
- Suppose $\exists S'$ with $S' \setminus A \neq \emptyset$ having $p_{S'}^{\mathbf{1}_A} > 0$. This would mean, for any $v \in S' \setminus A$, that $(\sum_S p_S^{\mathbf{1}_A} \mathbf{1}_S)(v) > 0$, a contradiction.
- Suppose $\exists S'$ s.t. $A \setminus S' \neq \emptyset$ with $p_{S'}^{\mathbf{1}_A} > 0$.
- Then, for any $v \in A \setminus S'$, consider below leading to a contradiction

$$\underbrace{p_{S'} \mathbf{1}_{S'}}_{>0} + \underbrace{\sum_{\substack{S \subseteq A \\ S \neq S'}} p_S \mathbf{1}_S}_{\text{can't sum to 1}} \Rightarrow \left(\sum_{\substack{S \subseteq A \\ S \neq S'}} p_S \mathbf{1}_S \right)(v) < 1 \quad (15.2)$$

$$1_{S'}(v) < 0$$

$$1_A(v) = 1$$

I.e., $v \in A$ so it must get value 1, but since $v \notin S'$, v is deficient. \square

Convexity of the Convex Closure

Lemma 15.3.2

$\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ is convex in $[0, 1]^V$.

Proof.

- Let $x, y \in [0, 1]^V$, $0 \leq \lambda \leq 1$, and $z = \lambda x + (1 - \lambda)y$, then

$$\lambda \check{f}(x) + (1 - \lambda) \check{f}(y) = \lambda \sum_S p_S^x f(S) + (1 - \lambda) \sum_S p_S^y f(S) \quad (15.3)$$

$$= \sum_S (\lambda p_S^x + (1 - \lambda) p_S^y) f(S) \quad (15.4)$$

$$= \sum_S p_S^{z'} f(S) \geq \min_{p \in \Delta^n(z)} E_{S \sim p}[f(S)] \quad (15.5)$$

$$= \check{f}(z) = \check{f}(\lambda x + (1 - \lambda)y) \quad (15.6)$$

$$p_S^{z'} \geq 0$$

$$\sum_S p_S^{z'} = 1$$

$$\sum_S p_S^{z'} \cdot 1_S = z$$

$$\therefore p^{z'} \in \Delta^n(z)$$

Convexity of the Convex Closure

Lemma 15.3.2

$\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ is convex in $[0, 1]^V$.

Proof.

- Let $x, y \in [0, 1]^V$, $0 \leq \lambda \leq 1$, and $z = \lambda x + (1 - \lambda)y$, then

$$\lambda \check{f}(x) + (1 - \lambda) \check{f}(y) = \lambda \sum_S p_S^x f(S) + (1 - \lambda) \sum_S p_S^y f(S) \quad (15.3)$$

$$= \sum_S (\lambda p_S^x + (1 - \lambda) p_S^y) f(S) \quad (15.4)$$

$$= \sum_S p_S^{z'} f(S) \geq \min_{p \in \Delta^n(z)} E_{S \sim p}[f(S)] \quad (15.5)$$

$$= \check{f}(z) = \check{f}(\lambda x + (1 - \lambda)y) \quad (15.6)$$

- Note that $p_S^{z'} = \lambda p_S^x + (1 - \lambda) p_S^y$ and is feasible in the min since $\sum_S p_S^{z'} = 1$, $p_S^{z'} \geq 0$ and $\sum_S p_S^{z'} \mathbf{1}_S = z$.

Def: Convex Envelope of a function

- Given any function $h : \mathbb{R}^n \rightarrow \mathbb{R}$, define new function $\check{h} : \mathbb{R}^n \rightarrow \mathbb{R}$ via:

$$\check{h}(x) = \sup \{g(x) : g \text{ is convex \& } g(y) \leq h(y), \forall y \in \mathbb{R}^n\} \quad (15.6)$$

Convex Closure is the Convex Envelope

Lemma 15.3.3

$\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ is the convex envelope.

Proof.

- Suppose \exists a convex \bar{f} with $\bar{f}(\mathbf{1}_A) = f(A) = \check{f}(\mathbf{1}_A), \forall A \subseteq V$ and $\exists x \in [0, 1]^V$ s.t. $\bar{f}(x) > \check{f}(x)$.
- Define p^x to be an achieving argmin in $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$. Hence, we have $x = \sum_S p_S^x \mathbf{1}_S$. Thus


$$\check{f}(x) = \sum_S p_S^x f(S) = \sum_S p_S^x \bar{f}(\mathbf{1}_S) \quad (15.7)$$

$$< \bar{f}(x) = \bar{f}\left(\sum_S p_S^x \mathbf{1}_S\right) \quad (15.8)$$

but this contradicts the convexity of \bar{f} .

Polymatroid with labeled edge lengths

- Recall $f(e|A) = f(A+e) - f(A)$
- Notice how submodularity, $f(e|B) \leq f(e|A)$ for $A \subseteq B$, defines the shape of the polytope.
- In fact, we have strictness here $f(e|B) < f(e|A)$ for $A \subset B$.
- Also, consider how the greedy algorithm proceeds along the edges of the polytope.




Polymatroid with labeled edge lengths

- Recall

$$f(e|A) = f(A+e) - f(A)$$
- Notice how submodularity,

$$f(e|B) \leq f(e|A) \text{ for } A \subseteq B,$$
 defines the shape of the polytope.
- In fact, we have strictness here

$$f(e|B) < f(e|A) \text{ for } A \subset B.$$
- Also, consider how the greedy algorithm proceeds along the edges of the polytope.



Optimization over P_f

- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

Optimization over P_f

- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

- Since P_f is down closed, if $\exists e \in E$ with $w(e) < 0$ then the solution above is unboundedly large.

Optimization over P_f

- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

- Since P_f is down closed, if $\exists e \in E$ with $w(e) < 0$ then the solution above is unboundedly large. Hence, assume $w \in \mathbb{R}_+^E$.

Optimization over P_f


- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

- Since P_f is down closed, if $\exists e \in E$ with $w(e) < 0$ then the solution above is unboundedly large. Hence, assume $w \in \mathbb{R}_+^E$.
- Due to **Theorem ??**, any $x \in P_f$ with $x \notin B_f$ is dominated by $x \leq y \in B_f$ which can only increase $w^\top x \leq w^\top y$ when $w \in \mathbb{R}_+^E$.

$x(u) \leq y(v)$ for.



Optimization over P_f

- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

- Since P_f is down closed, if $\exists e \in E$ with $w(e) < 0$ then the solution above is unboundedly large. Hence, assume $w \in \mathbb{R}_+^E$.
- Due to Theorem ??, any $x \in P_f$ with $x \notin B_f$ is dominated by $x \leq y \in B_f$ which can only increase $w^\top x \leq w^\top y$ when $w \in \mathbb{R}_+^E$.
- Hence, the problem is equivalent to: given $w \in \mathbb{R}_+^E$,

$$\text{maximize} \quad w^\top x \quad (15.10a)$$

$$\text{subject to} \quad x \in B_f \quad (15.10b)$$

Optimization over P_f

- Consider the following optimization. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.9a)$$

$$\text{subject to} \quad x \in P_f \quad (15.9b)$$

- Since P_f is down closed, if $\exists e \in E$ with $w(e) < 0$ then the solution above is unboundedly large. Hence, assume $w \in \mathbb{R}_+^E$.
- Due to Theorem ??, any $x \in P_f$ with $x \notin B_f$ is dominated by $x \leq y \in B_f$ which can only increase $w^\top x \leq w^\top y$ when $w \in \mathbb{R}_+^E$.
- Hence, the problem is equivalent to: given $w \in \mathbb{R}_+^E$,

$$\text{maximize} \quad w^\top x \quad (15.10a)$$

$$\text{subject to} \quad x \in B_f \quad (15.10b)$$

- Moreover, we can have $w \in \mathbb{R}^E$ if we insist on $x \in B_f$.

A continuous extension of f

- Consider again optimization problem. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.11a)$$

$$\text{subject to} \quad x \in B_f \quad (15.11b)$$

A continuous extension of f

- Consider again optimization problem. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.11a)$$

$$\text{subject to} \quad x \in B_f \quad (15.11b)$$

- We may consider this optimization problem a function $\check{f} : \mathbb{R}^E \rightarrow \mathbb{R}$ of $w \in \mathbb{R}^E$, defined as:

$$\check{f}(w) = \max\{wx : x \in B_f\} \quad (15.12)$$

A continuous extension of f

- Consider again optimization problem. Given $w \in \mathbb{R}^E$,

$$\text{maximize} \quad w^\top x \quad (15.11a)$$

$$\text{subject to} \quad x \in B_f \quad (15.11b)$$

- We may consider this optimization problem a function $\check{f} : \mathbb{R}^E \rightarrow \mathbb{R}$ of $w \in \mathbb{R}^E$, defined as:

$$\check{f}(w) = \max(w x : x \in B_f) \quad (15.12)$$

$\therefore \check{f}(w)$ is convex in w , max over lin. functions $w \cdot x$

- Hence, for any w , from the solution to the above theorem (as we have seen), we can compute the value of this function using Edmond's greedy algorithm.

Edmond's Theorem: The Greedy Algorithm

- Edmonds proved that the solution to $\check{f}(w) = \max\{wx : x \in B_f\}$ is solved by the greedy algorithm iff f is submodular.
- In particular, sort choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$.
- Define a vector $x^* \in \mathbb{R}^V$ where element e_i has value $x(e_i) = f(e_i | E_{i-1})$ for all $i \in V$.
- Then $\langle w, x^* \rangle = \max\{wx : x \in B_f\}$

Also true if
 1. change P_f to B_f
 2. $w \in \mathbb{R}^E$
 3. f is only submodular

Theorem 15.4.1 (Edmonds)

If $f : 2^E \rightarrow \mathbb{R}_+$ is given, and B is a polytope in \mathbb{R}_+^E of the form $B = \{x \in \mathbb{R}_+^E : x(A) \leq f(A), \forall A \subseteq E, x(E) = f(E)\}$, then the greedy solution to the problem $\max\{w^\top x : x \in P\}$ is $\forall w$ optimum iff f is monotone non-decreasing submodular (i.e., iff P is a polymatroid).

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w)$$

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.13)$$

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w) = \max\{wx : x \in B_f\} \quad (15.13)$$

$$= \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) = \sum_{i=1}^m w(e_i) x(e_i) \quad (15.14)$$

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w) = \max\{wx : x \in B_f\} \quad (15.13)$$

$$= \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) = \sum_{i=1}^m w(e_i) x(e_i) \quad (15.14)$$

$$= \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.15)$$

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w) = \max\{wx : x \in B_f\} \quad (15.13)$$

$$= \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) = \sum_{i=1}^m w(e_i) x(e_i) \quad (15.14)$$

$$= \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.15)$$

$$= w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \quad (15.16)$$

A continuous extension of submodular f

- That is, given a submodular function f , a $w \in \mathbb{R}^E$, choose element order (e_1, e_2, \dots, e_m) based on decreasing w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.
- Define the chain with i^{th} element $E_i = \{e_1, e_2, \dots, e_i\}$, we have

$$\check{f}(w) = \max\{wx : x \in B_f\} \quad (15.13)$$

$$= \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) = \sum_{i=1}^m w(e_i) x(e_i) \quad (15.14)$$

$$= \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.15)$$

$$= w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \quad (15.16)$$

- We say that $\emptyset \triangleq E_0 \subset E_1 \subset E_2 \subset \dots \subset E_m = E$ forms a **chain** based on w .

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

- Therefore, if f is a submodular function, we can write

$$\check{f}(w)$$

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

- Therefore, if f is a submodular function, we can write

$$\check{f}(w) = w(e_m)f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1}))f(E_i) \quad (15.18)$$

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

- Therefore, if f is a submodular function, we can write

$$\check{f}(w) = w(e_m)f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1}))f(E_i) \quad (15.18)$$

$$= \sum_{i=1}^m \lambda_i f(E_i) \quad (15.19)$$

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

- Therefore, if f is a submodular function, we can write

$$\check{f}(w) = w(e_m)f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1}))f(E_i) \quad (15.18)$$

$$= \sum_{i=1}^m \lambda_i f(E_i) \quad \lambda_i \geq 0 \quad w(e_i) \geq w(e_{i-1}) \quad (15.19)$$

where $\lambda_m = w(e_m)$ and otherwise $\lambda_i = w(e_i) - w(e_{i+1})$, where the elements are sorted descending according to w as before.

A continuous extension of submodular f

- Definition of the continuous extension, once again, for reference:

$$\check{f}(w) = \max(wx : x \in B_f) \quad (15.17)$$

- Therefore, if f is a submodular function, we can write

$$\check{f}(w) = w(e_m)f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1}))f(E_i) \quad (15.18)$$

$$= \sum_{i=1}^m \lambda_i f(E_i) \quad (15.19)$$

where $\lambda_m = w(e_m)$ and otherwise $\lambda_i = w(e_i) - w(e_{i+1})$, where the elements are sorted descending according to w as before.

- Convex analysis $\Rightarrow \check{f}(w) = \max(wx : x \in P)$ is always convex in w for any set $P \subseteq \mathbb{R}^E$, since a maximum of a set of linear functions (true even when f is not submodular or P is not itself a convex set).

An extension of f

- Recall, for any such $w \in \mathbb{R}^E$, we have

$$\begin{aligned}
 \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix} &= \underbrace{(w_1 - w_2)}_{\lambda_1} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \underbrace{(w_2 - w_3)}_{\lambda_2} \begin{pmatrix} 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \\
 &\cdots + \underbrace{(w_{n-1} - w_n)}_{\lambda_{m-1}} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \end{pmatrix} + \underbrace{(w_m)}_{\lambda_m} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{pmatrix} \tag{15.20}
 \end{aligned}$$

An extension of f

- Recall, for any such $w \in \mathbb{R}^E$, we have

$$\begin{aligned}
 \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix} &= \underbrace{(w_1 - w_2)}_{\lambda_1} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \underbrace{(w_2 - w_3)}_{\lambda_2} \begin{pmatrix} 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \\
 &\cdots + \underbrace{(w_{n-1} - w_n)}_{\lambda_{m-1}} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \end{pmatrix} + \underbrace{(w_m)}_{\lambda_m} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{pmatrix} \quad (15.20)
 \end{aligned}$$

- If we take w in decreasing order, then each coefficient of the vectors is non-negative (except possibly the last one, $\lambda_m = w_m$).

An extension of f

- Recall, for any such $w \in \mathbb{R}^E$, we have

$$\begin{aligned}
 \begin{pmatrix} w_1 \\ w_2 \\ \vdots \\ w_n \end{pmatrix} &= \underbrace{(w_1 - w_2)}_{\lambda_1} \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \underbrace{(w_2 - w_3)}_{\lambda_2} \begin{pmatrix} 1 \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix} + \dots \\
 &\dots + \underbrace{(w_{n-1} - w_n)}_{\lambda_{m-1}} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \end{pmatrix} + \underbrace{(w_m)}_{\lambda_m} \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 1 \end{pmatrix} \tag{15.20}
 \end{aligned}$$

$w \in [0, 1]^V$
 1) $\lambda_i \geq 0$
 2) $\lambda_i \leq 1$
 3) $\sum_i \lambda_i = v_i - v_n$
 $\lambda_0 = 1$
 $\sum_{i=0}^n \lambda_i = 1$

- If we take w in decreasing order, then each coefficient of the vectors is non-negative (except possibly the last one, $\lambda_m = w_m$).
- Often, we take $w \in \mathbb{R}_+^V$ or even $w \in [0, 1]^V$, where $\lambda_m \geq 0$.

An extension of f

- Define sets E_i based on this decreasing order of w as follows, for $i = 0, \dots, n$

$$E_i \stackrel{\text{def}}{=} \{e_1, e_2, \dots, e_i\} \quad (15.21)$$

An extension of f

- Define sets E_i based on this decreasing order of w as follows, for $i = 0, \dots, n$

$$E_i \stackrel{\text{def}}{=} \{e_1, e_2, \dots, e_i\} \quad (15.21)$$

- Note that

$$\mathbf{1}_{E_0} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \mathbf{1}_{E_1} = \begin{pmatrix} 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \dots, \mathbf{1}_{E_\ell} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \text{ etc.}$$

$\left. \begin{matrix} \vdots \\ \vdots \\ \vdots \\ \vdots \end{matrix} \right\} \ell \times$
 $\left. \begin{matrix} 0 \\ 0 \\ \vdots \\ 0 \end{matrix} \right\} (n - \ell) \times$

An extension of f

- Define sets E_i based on this decreasing order of w as follows, for $i = 0, \dots, n$

$$E_i \stackrel{\text{def}}{=} \{e_1, e_2, \dots, e_i\} \quad (15.21)$$

- Note that

$$\mathbf{1}_{E_0} = \begin{pmatrix} 0 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \mathbf{1}_{E_1} = \begin{pmatrix} 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \dots, \mathbf{1}_{E_\ell} = \begin{pmatrix} 1 \\ 1 \\ \vdots \\ 1 \\ 0 \\ \vdots \\ 0 \end{pmatrix}, \text{ etc.}$$

$\left. \begin{matrix} \vdots \\ \vdots \\ \vdots \\ \vdots \end{matrix} \right\} \ell \times$
 $\left. \begin{matrix} 0 \\ 0 \\ \vdots \\ 0 \end{matrix} \right\} (n - \ell) \times$

- Hence, from the previous and current slide, we have $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $1_A(i) = 1$ if $i \leq |A|$, and $1_A(i) = 0$ otherwise.

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $1_A(i) = 1$ if $i \leq |A|$, and $1_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\check{f}(w) \Big|_{w = \mathbf{1}_A}$$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $1_A(i) = 1$ if $i \leq |A|$, and $1_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i)$$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $1_A(i) = 1$ if $i \leq |A|$, and $1_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) = w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i)$$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $\mathbf{1}_A(i) = 1$ if $i \leq |A|$, and $\mathbf{1}_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\begin{aligned} \check{f}(w) &= \sum_{i=1}^m \lambda_i f(E_i) = w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \\ &= \mathbf{1}_A(m) f(E_m) + \sum_{i=1}^{m-1} (\mathbf{1}_A(i) - \mathbf{1}_A(i+1)) f(E_i) \end{aligned} \quad (15.23)$$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

so that $\mathbf{1}_A(i) = 1$ if $i \leq |A|$, and $\mathbf{1}_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\begin{aligned} \check{f}(w) &= \sum_{i=1}^m \lambda_i f(E_i) = w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \\ &= \mathbf{1}_A(m) f(E_m) + \sum_{i=1}^{m-1} (\mathbf{1}_A(i) - \mathbf{1}_A(i+1)) f(E_i) \quad (15.23) \\ &= (\underbrace{\mathbf{1}_A(|A|)}_{=1} - \underbrace{\mathbf{1}_A(|A|+1)}_{=0}) f(E_{|A|}) = f(E_{|A|}) \end{aligned}$$

From \check{f} back to f , even when f is not submodular

- From the continuous \check{f} , we can recover $f(A)$ for any $A \subseteq V$.
- Take $w = \mathbf{1}_A$ for some $A \subseteq E$, so w is vertex of the hypercube.
- Order the elements of E in decreasing order of w so that $w(e_1) \geq w(e_2) \geq w(e_3) \geq \dots \geq w(e_m)$.
- This means

$$w = (w(e_1), w(e_2), \dots, w(e_m)) = (\underbrace{1, 1, 1, \dots, 1}_{|A| \text{ times}}, \underbrace{0, 0, \dots, 0}_{m-|A| \text{ times}}) \quad (15.22)$$

assuming element names are so ordered.

so that $\mathbf{1}_A(i) = 1$ if $i \leq |A|$, and $\mathbf{1}_A(i) = 0$ otherwise.

- For any $f : 2^E \rightarrow \mathbb{R}$, $w = \mathbf{1}_A$, since $E_{|A|} = \{e_1, e_2, \dots, e_{|A|}\} = A$:

$$\begin{aligned} \check{f}(w) &= \sum_{i=1}^m \lambda_i f(E_i) = w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \\ &= \mathbf{1}_A(m) f(E_m) + \sum_{i=1}^{m-1} (\mathbf{1}_A(i) - \mathbf{1}_A(i+1)) f(E_i) \end{aligned} \quad (15.23)$$

$$= (\mathbf{1}_A(|A|) - \mathbf{1}_A(|A| + 1)) f(E_{|A|}) = f(E_{|A|}) = f(A) \quad (15.24)$$

From \check{f} back to f

- We can view $\check{f} : [0, 1]^E \rightarrow \mathbb{R}$ defined on the hypercube, with f defined as \check{f} evaluated on the hypercube extreme points (vertices).

From \check{f} back to f

- We can view $\check{f} : [0, 1]^E \rightarrow \mathbb{R}$ defined on the hypercube, with f defined as \check{f} evaluated on the hypercube extreme points (vertices).
- To summarize, with $\check{f}(\mathbf{1}_A) = \sum_{i=1}^m \lambda_i f(E_i)$, we have

$$\check{f}(\mathbf{1}_A) = f(A), \quad (15.25)$$

From \check{f} back to f

- We can view $\check{f} : [0, 1]^E \rightarrow \mathbb{R}$ defined on the hypercube, with f defined as \check{f} evaluated on the hypercube extreme points (vertices).
- To summarize, with $\check{f}(\mathbf{1}_A) = \sum_{i=1}^m \lambda_i f(E_i)$, we have

$$\check{f}(\mathbf{1}_A) = f(A), \quad (15.25)$$

- ... and when f is submodular, we also have have

$$\check{f}(\mathbf{1}_A) = \max \{ \mathbf{1}_A^\top x : x \in B_f \} \quad (15.26)$$

$$= \max \{ \mathbf{1}_A^\top x : x(B) \leq f(B), \forall B \subseteq E \} \quad (15.27)$$

From \check{f} back to f

- We can view $\check{f} : [0, 1]^E \rightarrow \mathbb{R}$ defined on the hypercube, with f defined as \check{f} evaluated on the hypercube extreme points (vertices).
- To summarize, with $\check{f}(\mathbf{1}_A) = \sum_{i=1}^m \lambda_i f(E_i)$, we have

$$\check{f}(\mathbf{1}_A) = f(A), \quad (15.25)$$

- ... and when f is submodular, we also have have

$$\check{f}(\mathbf{1}_A) = \max \{ \mathbf{1}_A^\top x : x \in B_f \} \quad (15.26)$$

$$= \max \{ \mathbf{1}_A^\top x : x(B) \leq f(B), \forall B \subseteq E \} \quad (15.27)$$

- Note when considering only $\check{f} : [0, 1]^E \rightarrow \mathbb{R}$, then any $w \in [0, 1]^E$ is in positive orthant, and we have

$$\check{f}(w) = \max \{ w^\top x : x \in P_f \} \quad (15.28)$$

An extension of an arbitrary $f : 2^V \rightarrow \mathbb{R}$

- Thus, for any $f : 2^E \rightarrow \mathbb{R}$, even non-submodular f , we can define an extension, having $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, in this way where

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.29)$$

with the $E_i = \{e_1, \dots, e_i\}$'s defined based on sorted descending order of w as in $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$, and where

$$\text{for } i \in \{1, \dots, m\}, \quad \lambda_i = \begin{cases} w(e_i) - w(e_{i+1}) & \text{if } i < m \\ w(e_m) & \text{if } i = m \end{cases} \quad (15.30)$$

so that $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$.

An extension of an arbitrary $f : 2^V \rightarrow \mathbb{R}$

- Thus, for any $f : 2^E \rightarrow \mathbb{R}$, even non-submodular f , we can define an extension, having $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, in this way where

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.29)$$

with the $E_i = \{e_1, \dots, e_i\}$'s defined based on sorted descending order of w as in $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$, and where

$$\text{for } i \in \{1, \dots, m\}, \quad \lambda_i = \begin{cases} w(e_i) - w(e_{i+1}) & \text{if } i < m \\ w(e_m) & \text{if } i = m \end{cases} \quad (15.30)$$

so that $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$.

- $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ is an interpolation of certain hypercube vertices.

An extension of an arbitrary $f : 2^V \rightarrow \mathbb{R}$

- Thus, for any $f : 2^E \rightarrow \mathbb{R}$, even non-submodular f , we can define an extension, having $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, in this way where

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.29)$$

with the $E_i = \{e_1, \dots, e_i\}$'s defined based on sorted descending order of w as in $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$, and where

$$\text{for } i \in \{1, \dots, m\}, \quad \lambda_i = \begin{cases} w(e_i) - w(e_{i+1}) & \text{if } i < m \\ w(e_m) & \text{if } i = m \end{cases} \quad (15.30)$$

so that $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$.

- $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ is an interpolation of certain hypercube vertices.
- $\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i)$ is the associated interpolation of the values of f at sets corresponding to each hypercube vertex.

An extension of an arbitrary $f : 2^V \rightarrow \mathbb{R}$

- Thus, for any $f : 2^E \rightarrow \mathbb{R}$, even non-submodular f , we can define an extension, having $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, in this way where

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.29)$$

with the $E_i = \{e_1, \dots, e_i\}$'s defined based on sorted descending order of w as in $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$, and where

$$\text{for } i \in \{1, \dots, m\}, \quad \lambda_i = \begin{cases} w(e_i) - w(e_{i+1}) & \text{if } i < m \\ w(e_m) & \text{if } i = m \end{cases} \quad (15.30)$$

so that $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$.

- $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ is an interpolation of certain hypercube vertices.
- $\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i)$ is the associated interpolation of the values of f at sets corresponding to each hypercube vertex.
- This extension is called the Lovász extension!

Weighted gains vs. weighted functions

- Again sorting E descending in w , the extension summarized:

$$\check{f}(w) = \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) \quad (15.31)$$

$$= \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.32)$$

$$= w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \quad (15.33)$$

$$= \sum_{i=1}^m \lambda_i f(E_i) \quad (15.34)$$

Weighted gains vs. weighted functions

- Again sorting E descending in w , the extension summarized:

$$\check{f}(w) = \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) \quad \text{currently assumed } f(\emptyset) = 0. \quad (15.31)$$

$$= \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.32)$$

$$= w(e_m) f(E_m) + \sum_{i=1}^{m-1} (w(e_i) - w(e_{i+1})) f(E_i) \quad (15.33)$$

$$= \sum_{i=1}^m \lambda_i f(E_i) \quad (15.34)$$

- So $\check{f}(w)$ seen either as **sum of weighted gain evaluations** (Eqn. (15.31)), or as **sum of weighted function evaluations** (Eqn. (15.34)).

Summary: comparison of the two extension forms

- So if f is **submodular**, then we can write $\check{f}(w) = \max\{wx : x \in B_f\}$ (which is clearly convex) in the form:

$$\check{f}(w) = \max\{wx : x \in B_f\} = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.35)$$

from Edment gradg.

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

Summary: comparison of the two extension forms

- So if f is **submodular**, then we can write $\check{f}(w) = \max(wx : x \in B_f)$ (which is clearly convex) in the form:

$$\check{f}(w) = \max(wx : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.35)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

- On the other hand, for any f (even non-submodular), we can produce an extension \check{f} having the form

so right

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.36)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

called the L. E.

Summary: comparison of the two extension forms

- So if f is **submodular**, then we can write $\check{f}(w) = \max(wx : x \in B_f)$ (which is clearly convex) in the form:

$$\check{f}(w) = \max(wx : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.35)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

- On the other hand, **for any f (even non-submodular)**, we can produce an extension \check{f} having the form

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.36)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

- In both Eq. (15.35) and Eq. (15.36), we have $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, but Eq. (15.36), might not be convex.

Summary: comparison of the two extension forms

- So if f is **submodular**, then we can write $\check{f}(w) = \max(wx : x \in B_f)$ (which is clearly convex) in the form:

$$\check{f}(w) = \max(wx : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.35)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

- On the other hand, for any f (even non-submodular), we can produce an extension \check{f} having the form

$$\check{f}(w) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.36)$$

where $w = \sum_{i=1}^m \lambda_i \mathbf{1}_{E_i}$ and $E_i = \{e_1, \dots, e_i\}$ defined based on sorted descending order $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

- In both Eq. (15.35) and Eq. (15.36), we have $\check{f}(\mathbf{1}_A) = f(A)$, $\forall A$, but Eq. (15.36), might not be convex.
- Submodularity is sufficient for convexity, but is it necessary?

The Lovász extension of $f : 2^E \rightarrow \mathbb{R}$

- Lovász showed that if a function $\check{f}(w)$ defined as in Eqn. (15.29) is convex, then f must be submodular.

15.36

(15.29)

The Lovász extension of $f : 2^E \rightarrow \mathbb{R}$

- Lovász showed that if a function $\check{f}(w)$ defined as in Eqn. (15.29) is convex, then f must be submodular.
- This **continuous extension** \check{f} of f , in any case (f being submodular or not), is typically called the **Lovász extension** of f (but also sometimes called the Choquet integral, or the Lovász-Edmonds extension).

Lovász Extension, Submodularity and Convexity

Theorem 15.4.2

A function $f : 2^E \rightarrow \mathbb{R}$ is submodular iff its Lovász extension \check{f} of f is convex.

Proof.

- We've already seen that if f is submodular, its extension can be written via Eqn.(15.29) due to the greedy algorithm, and therefore is also equivalent to $\check{f}(w) = \max \{wx : x \in P_f\}$, and thus is convex.

Lovász Extension, Submodularity and Convexity

Theorem 15.4.2

A function $f : 2^E \rightarrow \mathbb{R}$ is submodular iff its Lovász extension \check{f} of f is convex.

Proof.

- We've already seen that if f is submodular, its extension can be written via Eqn.(15.29) due to the greedy algorithm, and therefore is also equivalent to $\check{f}(w) = \max \{wx : x \in P_f\}$, and thus is convex.
- Conversely, suppose the Lovász extension $\check{f}(w) = \sum_i \lambda_i f(E_i)$ of some function $f : 2^E \rightarrow \mathbb{R}$ is a convex function.

...

Lovász Extension, Submodularity and Convexity

Theorem 15.4.2

A function $f : 2^E \rightarrow \mathbb{R}$ is submodular iff its Lovász extension \check{f} of f is convex.

Proof.

- We've already seen that if f is submodular, its extension can be written via Eqn.(15.29) due to the greedy algorithm, and therefore is also equivalent to $\check{f}(w) = \max \{wx : x \in P_f\}$, and thus is convex.
- Conversely, suppose the Lovász extension $\check{f}(w) = \sum_i \lambda_i f(E_i)$ of some function $f : 2^E \rightarrow \mathbb{R}$ is a convex function.
- We note that, based on the extension definition, in particular the definition of the $\{\lambda_i\}_i$, we have that $\check{f}(\alpha w) = \alpha \check{f}(w)$ for any $\alpha \in \mathbb{R}_+$. I.e., f is a positively homogeneous convex function.

...

Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Earlier, we saw that $\check{f}(\mathbf{1}_A) = f(A)$ for all $A \subseteq E$. \Rightarrow tight.

Lovász Extension, Submodularity and Convexity


... proof of Thm. 15.4.2 cont.

- Earlier, we saw that $\check{f}(\mathbf{1}_A) = f(A)$ for all $A \subseteq E$.
- Now, given $A, B \subseteq E$, we will show that

$$\check{f}(\mathbf{1}_A + \mathbf{1}_B) = \check{f}(\mathbf{1}_{A \cup B} + \mathbf{1}_{A \cap B}) \quad (15.37)$$

$$= f(A \cup B) + f(A \cap B). \quad (15.38)$$

$$(\mathbf{1}_A + \mathbf{1}_B)(v) = \begin{cases} 2 & \text{if } v \in A \cap B \\ 1 & \text{if } v \in A \Delta B \\ 0 & \text{else.} \end{cases}$$



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Earlier, we saw that $\check{f}(\mathbf{1}_A) = f(A)$ for all $A \subseteq E$.
- Now, given $A, B \subseteq E$, we will show that

$$\check{f}(\mathbf{1}_A + \mathbf{1}_B) = \check{f}(\mathbf{1}_{A \cup B} + \mathbf{1}_{A \cap B}) \quad (15.37)$$

$$= f(A \cup B) + f(A \cap B). \quad (15.38)$$

- Let $C = A \cap B$, order E based on decreasing $w = \mathbf{1}_A + \mathbf{1}_B$ so that

$$w = (w(e_1), w(e_2), \dots, w(e_m)) \quad (15.39)$$

$$= \underbrace{(2, 2, \dots, 2)}_{i \in C}, \underbrace{(1, 1, \dots, 1)}_{i \in A \Delta B}, \underbrace{(0, 0, \dots, 0)}_{i \in E \setminus (A \cup B)} \quad (15.40)$$

Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Earlier, we saw that $\check{f}(\mathbf{1}_A) = f(A)$ for all $A \subseteq E$.
- Now, given $A, B \subseteq E$, we will show that

$$\check{f}(\mathbf{1}_A + \mathbf{1}_B) = \check{f}(\mathbf{1}_{A \cup B} + \mathbf{1}_{A \cap B}) \quad (15.37)$$

$$= f(A \cup B) + f(A \cap B). \quad (15.38)$$

- Let $C = A \cap B$, order E based on decreasing $w = \mathbf{1}_A + \mathbf{1}_B$ so that

$$w = (w(e_1), w(e_2), \dots, w(e_m)) \quad (15.39)$$

$$= (\underbrace{2, 2, \dots, 2}_{i \in C}, \underbrace{1, 1, \dots, 1}_{i \in A \Delta B}, \underbrace{0, 0, \dots, 0}_{i \in E \setminus (A \cup B)}) \quad (15.40)$$

- Then, considering $\check{f}(w) = \sum_i \lambda_i f(E_i)$, we have $\lambda_{|C|} = 1$, $\lambda_{|A \cup B|} = 1$, and $\lambda_i = 0$ for $i \notin \{|C|, |A \cup B|\}$.

$\lambda_m = w_m$ also zero

Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Earlier, we saw that $\check{f}(\mathbf{1}_A) = f(A)$ for all $A \subseteq E$.
- Now, given $A, B \subseteq E$, we will show that

$$\check{f}(\mathbf{1}_A + \mathbf{1}_B) = \check{f}(\mathbf{1}_{A \cup B} + \mathbf{1}_{A \cap B}) \quad (15.37)$$

$$= f(A \cup B) + f(A \cap B). \quad (15.38)$$

- Let $C = A \cap B$, order E based on decreasing $w = \mathbf{1}_A + \mathbf{1}_B$ so that

$$w = (w(e_1), w(e_2), \dots, w(e_m)) \quad (15.39)$$

$$= (\underbrace{2, 2, \dots, 2}_{i \in C}, \underbrace{1, 1, \dots, 1}_{i \in A \Delta B}, \underbrace{0, 0, \dots, 0}_{i \in E \setminus (A \cup B)}) \quad (15.40)$$

- Then, considering $\check{f}(w) = \sum_i \lambda_i f(E_i)$, we have $\lambda_{|C|} = 1$, $\lambda_{|A \cup B|} = 1$, and $\lambda_i = 0$ for $i \notin \{|C|, |A \cup B|\}$.
- But then $E_{|C|} = A \cap B$ and $E_{|A \cup B|} = A \cup B$. Therefore, $\check{f}(w) = \check{f}(\mathbf{1}_A + \mathbf{1}_B) = f(A \cap B) + f(A \cup B)$.

Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)]$$

(15.44)



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)] = 0.5[\check{f}(\mathbf{1}_A + \mathbf{1}_B)] \quad (15.41)$$

(15.44)



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)] = 0.5[\check{f}(\mathbf{1}_A + \mathbf{1}_B)] \quad (15.41)$$

$$= \check{f}(0.5\mathbf{1}_A + 0.5\mathbf{1}_B) \quad (15.42)$$

$$(15.44)$$



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)] = 0.5[\check{f}(\mathbf{1}_A + \mathbf{1}_B)] \quad (15.41)$$

$$= \check{f}(0.5\mathbf{1}_A + 0.5\mathbf{1}_B) \quad (15.42)$$

$$\leq 0.5\check{f}(\mathbf{1}_A) + 0.5\check{f}(\mathbf{1}_B) \quad (15.43)$$

$$(15.44)$$



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)] = 0.5[\check{f}(\mathbf{1}_A + \mathbf{1}_B)] \quad (15.41)$$

$$= \check{f}(0.5\mathbf{1}_A + 0.5\mathbf{1}_B) \quad (15.42)$$

$$\leq 0.5\check{f}(\mathbf{1}_A) + 0.5\check{f}(\mathbf{1}_B) \quad (15.43)$$

$$= 0.5(f(A) + f(B)) \quad (15.44)$$



Lovász Extension, Submodularity and Convexity

... proof of Thm. 15.4.2 cont.

- Also, since \check{f} is convex (by assumption) and positively homogeneous, we have for any $A, B \subseteq E$,

$$0.5[f(A \cap B) + f(A \cup B)] = 0.5[\check{f}(\mathbf{1}_A + \mathbf{1}_B)] \stackrel{\check{f}(0.5\mathbf{1}_A) + \check{f}(0.5\mathbf{1}_B)}{=} \quad (15.41)$$

$$= \check{f}(0.5\mathbf{1}_A + 0.5\mathbf{1}_B) \quad (15.42)$$

$$\leq 0.5\check{f}(\mathbf{1}_A) + 0.5\check{f}(\mathbf{1}_B) \quad (15.43)$$

$$= 0.5(f(A) + f(B)) \quad (15.44)$$

- Thus, we have shown that for any $A, B \subseteq E$,

$$f(A \cup B) + f(A \cap B) \leq f(A) + f(B) \quad (15.45)$$

so f must be submodular.

$$x = 0.5 \cdot (2x)$$

Cont. Submodularity

$$f(x+y) \leq f(x) + f(y)$$

$$f(0.5x + 0.5y) \leq 0.5 \cdot f(x) + 0.5 \cdot f(y) = f(0.5x) + f(0.5y)$$

□

Lovász ext. vs. the concave closure of submodular function

- The above theorem showed that the Lovász extension is convex iff f is submodular.

Lovász ext. vs. the concave closure of submodular function

- The above theorem showed that the Lovász extension is convex iff f is submodular.
- Our next theorem shows that the Lovász extension coincides precisely with the convex closure iff f is submodular.

Lovász ext. vs. the concave closure of submodular function

- The above theorem showed that the Lovász extension is convex iff f is submodular.
- Our next theorem shows that the Lovász extension coincides precisely with the convex closure iff f is submodular.
- I.e., not only is the Lovász extension convex for f submodular, it is the convex closure when f is convex.

Lovász ext. vs. the concave closure of submodular function

- The above theorem showed that the Lovász extension is convex iff f is submodular.
- Our next theorem shows that the Lovász extension coincides precisely with the convex closure iff f is submodular.
- I.e., not only is the Lovász extension convex for f submodular, it is the convex closure when f is convex.
- Hence, convex closure is easy to evaluate when f is submodular and is this particular form iff f is submodular.

Lovász ext. vs. the concave closure of submodular function

Theorem 15.4.3

Let $\check{f}(w) = \max\{wx : x \in B_f\} = \sum_{i=1}^m \lambda_i f(E_i)$ be the Lovász extension and $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ be the convex closure. Then \check{f} and \check{f} coincide iff f is submodular.

from only if f submodular.

Proof.

- Assume f is submodular.

Lovász ext. vs. the concave closure of submodular function

Theorem 15.4.3

Let $\check{f}(w) = \max(w x : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i)$ be the Lovász extension and $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ be the convex closure. Then \check{f} and \check{f} coincide iff f is submodular.

Proof.

- Assume f is submodular.
- Given x , let p^x be an achieving argmin in $\check{f}(x)$ that also maximizes $\sum_S p_S^x |S|^2$.


Lovász ext. vs. the concave closure of submodular function

Theorem 15.4.3

Let $\check{f}(w) = \max(w x : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i)$ be the Lovász extension and $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ be the convex closure. Then \check{f} and \check{f} coincide iff f is submodular.

Proof.

- Assume f is submodular.
- Given x , let p^x be an achieving argmin in $\check{f}(x)$ that also maximizes $\sum_S p_S^x |S|^2$.
- Suppose $\exists A, B \subseteq V$ that are crossing (i.e., $A \not\subseteq B, B \not\subseteq A$) and positive and w.l.o.g., $\underline{p_A^x} \geq p_B^x > 0$.



Lovász ext. vs. the concave closure of submodular function

Theorem 15.4.3

Let $\check{f}(w) = \max(w x : x \in B_f) = \sum_{i=1}^m \lambda_i f(E_i)$ be the Lovász extension and $\check{f}(x) = \min_{p \in \Delta^n(x)} E_{S \sim p}[f(S)]$ be the convex closure. Then \check{f} and \check{f} coincide iff f is submodular.

Proof.

- Assume f is submodular.
- Given x , let p^x be an achieving argmin in $\check{f}(x)$ that also maximizes $\sum_S p_S^x |S|^2$.
- Suppose $\exists A, B \subseteq V$ that are crossing (i.e., $A \not\subseteq B, B \not\subseteq A$) and positive and w.l.o.g., $p_A^x \geq p_B^x > 0$.
- Then we may update p^x as follows:

$$\bar{p}_A^x \leftarrow p_A^x - p_B^x \qquad \bar{p}_B^x \leftarrow p_B^x - p_B^x \qquad (15.46)$$

$$\bar{p}_{A \cup B}^x \leftarrow p_{A \cup B}^x + p_B^x \qquad \bar{p}_{A \cap B}^x \leftarrow p_{A \cap B}^x + p_B^x \qquad (15.47)$$

and by submodularity, this does not increase $\sum_S p_S^x f(S)$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- This does increase $\sum_S p_S^x |S|^2$ however since

$$|A \cup B|^2 + |A \cap B|^2 = (|A| + |B \setminus A|)^2 + (|B| - |B \setminus A|)^2 \quad (15.48)$$

$$= |A|^2 + |B|^2 + 2|B \setminus A|(|A| - |B| + |B \setminus A|) \quad (15.49)$$

$$\geq |A|^2 + |B|^2 \quad \geq 0 \quad (15.50)$$

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- This does increase $\sum_S p_S^x |S|^2$ however since

$$|A \cup B|^2 + |A \cap B|^2 = (|A| + |B \setminus A|)^2 + (|B| - |B \setminus A|)^2 \quad (15.48)$$

$$= |A|^2 + |B|^2 + 2|B \setminus A|(|A| - |B| + |B \setminus A|) \quad (15.49)$$

$$\overset{>}{\text{red}} |A|^2 + |B|^2 \quad \overset{>0}{\text{red}} \quad \overset{>0}{\text{red}} \quad (15.50)$$

- Contradiction! Hence, there can be no crossing sets A, B and we must have, for any A, B with $p_A^x > 0$ and $p_B^x > 0$ either $A \subset B$ or $B \subset A$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- This does increase $\sum_S p_S^x |S|^2$ however since

$$|A \cup B|^2 + |A \cap B|^2 = (|A| + |B \setminus A|)^2 + (|B| - |B \setminus A|)^2 \quad (15.48)$$

$$= |A|^2 + |B|^2 + 2|B \setminus A|(|A| - |B| + |B \setminus A|) \quad (15.49)$$

$$\geq |A|^2 + |B|^2 \quad (15.50)$$

- Contradiction! Hence, there can be no crossing sets A, B and we must have, for any A, B with $p_A^x > 0$ and $p_B^x > 0$ either $A \subset B$ or $B \subset A$.
- Hence, the sets $\{A \subseteq V : p_A^x > 0\}$ form a chain and can be as large only as size $n = |V|$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- This does increase $\sum_S p_S^x |S|^2$ however since

$$|A \cup B|^2 + |A \cap B|^2 = (|A| + |B \setminus A|)^2 + (|B| - |B \setminus A|)^2 \quad (15.48)$$

$$= |A|^2 + |B|^2 + 2|B \setminus A|(|A| - |B| + |B \setminus A|) \quad (15.49)$$

$$\geq |A|^2 + |B|^2 \quad (15.50)$$

- Contradiction! Hence, there can be no crossing sets A, B and we must have, for any A, B with $p_A^x > 0$ and $p_B^x > 0$ either $A \subset B$ or $B \subset A$.
- Hence, the sets $\{A \subseteq V : p_A^x > 0\}$ form a chain and can be as large only as size $n = |V|$.
- This is the same chain that defines the Lovász extension $\check{f}(x)$, namely $\emptyset = E_0 \subseteq E_1 \subseteq E_2 \subseteq \dots$ where $E_i = \{e_1, e_2, \dots, e_i\}$ and e_i is ordered so that $x(e_1) \geq x(e_2) \geq \dots \geq x(e_n)$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.
- Since f is not submodular, $\exists S$ and $i, j \notin S$ such that $f(S) + f(S + i + j) > f(S + i) + f(S + j)$, a strict violation of submodularity.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.
- Since f is not submodular, $\exists S$ and $i, j \notin S$ such that $f(S) + f(S + i + j) > f(S + i) + f(S + j)$, a strict violation of submodularity.
- Consider $x = \mathbf{1}_S + \frac{1}{2}\mathbf{1}_{\{i,j\}}$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.
- Since f is not submodular, $\exists S$ and $i, j \notin S$ such that $f(S) + f(S + i + j) > f(S + i) + f(S + j)$, a strict violation of submodularity.
- Consider $x = \mathbf{1}_S + \frac{1}{2}\mathbf{1}_{\{i,j\}}$.
- Then $\check{f}(x) = \frac{1}{2}f(S) + \frac{1}{2}f(S + i + j)$ and p^x is feasible for \check{f} with $p_S^x = 1/2$ and $p_{S+i+j}^x = 1/2$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.
- Since f is not submodular, $\exists S$ and $i, j \notin S$ such that $f(S) + f(S + i + j) > f(S + i) + f(S + j)$, a strict violation of submodularity.
- Consider $x = \mathbf{1}_S + \frac{1}{2}\mathbf{1}_{\{i,j\}}$.
- Then $\check{f}(x) = \frac{1}{2}f(S) + \frac{1}{2}f(S + i + j)$ and p^x is feasible for \check{f} with $p_S^x = 1/2$ and $p_{S+i+j}^x = 1/2$.
- An alternate feasible distribution for x in the convex closure is $\bar{p}_{S+i}^x = \bar{p}_{S+j}^x = 1/2$.

Lovász ext. vs. the concave closure of submodular function

... proof cont.

- Next, assume f is not submodular. We must show that the Lovász extension $\check{f}(x)$ and the concave closure $\check{\check{f}}(x)$ need not coincide.
- Since f is not submodular, $\exists S$ and $i, j \notin S$ such that $f(S) + f(S + i + j) > f(S + i) + f(S + j)$, a strict violation of submodularity.
- Consider $x = \mathbf{1}_S + \frac{1}{2}\mathbf{1}_{\{i,j\}}$.
- Then $\check{f}(x) = \frac{1}{2}f(S) + \frac{1}{2}f(S + i + j)$ and p^x is feasible for \check{f} with $p_S^x = 1/2$ and $p_{S+i+j}^x = 1/2$.
- An alternate feasible distribution for x in the convex closure is $\bar{p}_{S+i}^x = \bar{p}_{S+j}^x = 1/2$.
- This gives

$$\check{\check{f}}(x) \leq \frac{1}{2}[f(S + i) + f(S + j)] < \check{f}(x) \quad (15.51)$$

meaning $\check{f}(x) \neq \check{\check{f}}(x)$.

Integration and Aggregation

- Integration is just summation (e.g., the \int symbol has as its origins a sum).

Integration and Aggregation

- Integration is just summation (e.g., the \int symbol has as its origins a sum).
- **Lebesgue integration** allows integration w.r.t. an underlying measure μ of sets. E.g., given measurable function f , we can define

$$\int_X f du = \sup I_X(s) \quad (15.52)$$

where $I_X(s) = \sum_{i=1}^n c_i \mu(X \cap X_i)$, and where we take the sup over all measurable functions s such that $0 \leq s \leq f$ and $s(x) = \sum_{i=1}^n c_i I_{X_i}(x)$ and where $I_{X_i}(x)$ is indicator of membership of set X_i , with $c_i > 0$.

Integration, Aggregation, and Weighted Averages

- In finite discrete spaces, Lebesgue integration is just a weighted average, and can be seen as an **aggregation** function.

Integration, Aggregation, and Weighted Averages

- In finite discrete spaces, Lebesgue integration is just a weighted average, and can be seen as an **aggregation** function.
- I.e., given a weight vector $w \in [0, 1]^E$ for some finite ground set E , then for any $x \in \mathbb{R}^E$ we have the weighted average of x as:

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.53)$$

Integration, Aggregation, and Weighted Averages

- In finite discrete spaces, Lebesgue integration is just a weighted average, and can be seen as an **aggregation** function.
- I.e., given a weight vector $w \in [0, 1]^E$ for some finite ground set E , then for any $x \in \mathbb{R}^E$ we have the weighted average of x as:

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.53)$$

- Consider $\mathbf{1}_e$ for $e \in E$, we have

$$\text{WAVG}(\mathbf{1}_e) = w(e) \quad (15.54)$$

Integration, Aggregation, and Weighted Averages

- In finite discrete spaces, Lebesgue integration is just a weighted average, and can be seen as an **aggregation** function.
- I.e., given a weight vector $w \in [0, 1]^E$ for some finite ground set E , then for any $x \in \mathbb{R}^E$ we have the weighted average of x as:

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.53)$$

- Consider $\mathbf{1}_e$ for $e \in E$, we have

$$\text{WAVG}(\mathbf{1}_e) = w(e) \quad (15.54)$$

so seen as a function on the hypercube vertices, the entire WAVG function is given based on values on a size $m = |E|$ **subset** of the vertices of this hypercube, i.e., $\{\mathbf{1}_e : e \in E\}$.

Integration, Aggregation, and Weighted Averages

- In finite discrete spaces, Lebesgue integration is just a weighted average, and can be seen as an **aggregation** function.
- I.e., given a weight vector $w \in [0, 1]^E$ for some finite ground set E , then for any $x \in \mathbb{R}^E$ we have the weighted average of x as:

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.53)$$

- Consider $\mathbf{1}_e$ for $e \in E$, we have

$$\text{WAVG}(\mathbf{1}_e) = w(e) \quad (15.54)$$

so seen as a function on the hypercube vertices, the entire WAVG function is given based on values on a size $m = |E|$ **subset** of the vertices of this hypercube, i.e., $\{\mathbf{1}_e : e \in E\}$. **Moreover, we are interpolating as in**

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) = \sum_{e \in E} x(e)\text{WAVG}(\mathbf{1}_e) \quad (15.55)$$

Integration, Aggregation, and Weighted Averages

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.56)$$

- Clearly, WAVG function is linear in weights w , in the argument x , and is homogeneous. That is, for all $w, w_1, w_2, x, x_1, x_2 \in \mathbb{R}^E$ and $\alpha \in \mathbb{R}$,

$$\text{WAVG}_{w_1+w_2}(x) = \text{WAVG}_{w_1}(x) + \text{WAVG}_{w_2}(x), \quad (15.57)$$

$$\text{WAVG}_w(x_1 + x_2) = \text{WAVG}_w(x_1) + \text{WAVG}_w(x_2), \quad (15.58)$$

and,

$$\text{WAVG}(\alpha x) = \alpha \text{WAVG}(x). \quad (15.59)$$

Integration, Aggregation, and Weighted Averages

$$\text{WAVG}(x) = \sum_{e \in E} x(e)w(e) \quad (15.56)$$

- Clearly, WAVG function is linear in weights w , in the argument x , and is homogeneous. That is, for all $w, w_1, w_2, x, x_1, x_2 \in \mathbb{R}^E$ and $\alpha \in \mathbb{R}$,

$$\text{WAVG}_{w_1+w_2}(x) = \text{WAVG}_{w_1}(x) + \text{WAVG}_{w_2}(x), \quad (15.57)$$

$$\text{WAVG}_w(x_1 + x_2) = \text{WAVG}_w(x_1) + \text{WAVG}_w(x_2), \quad (15.58)$$

and,

$$\text{WAVG}(\alpha x) = \alpha \text{WAVG}(x). \quad (15.59)$$

- We will see: The Lovász extension is still be linear in “weights” (i.e., the submodular function f), but will not be linear in x and will only be positively homogeneous (for $\alpha \geq 0$).

Integration, Aggregation, and Weighted Averages

- More complex “nonlinear” aggregation functions can be constructed by defining the aggregation function on all vertices of the hypercube. I.e., for each $\mathbf{1}_A : A \subseteq E$ we might have (for all $A \subseteq E$):

$$\text{AG}(\mathbf{1}_A) = w_A \quad (15.60)$$

Integration, Aggregation, and Weighted Averages

- More complex “nonlinear” aggregation functions can be constructed by defining the aggregation function on all vertices of the hypercube. I.e., for each $\mathbf{1}_A : A \subseteq E$ we might have (for all $A \subseteq E$):

$$\text{AG}(\mathbf{1}_A) = w_A \quad (15.60)$$

- What then might $\text{AG}(x)$ be for some $x \in \mathbb{R}^E$? Our weighted average functions might look **something** more like the r.h.s. in:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.61)$$

Integration, Aggregation, and Weighted Averages

- More complex “nonlinear” aggregation functions can be constructed by defining the aggregation function on all vertices of the hypercube. I.e., for each $\mathbf{1}_A : A \subseteq E$ we might have (for all $A \subseteq E$):

$$\text{AG}(\mathbf{1}_A) = w_A \quad (15.60)$$

- What then might $\text{AG}(x)$ be for some $x \in \mathbb{R}^E$? Our weighted average functions might look **something** more like the r.h.s. in:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.61)$$

- Note, we can define $w(e) = w'(e)$ and $w(A) = 0, \forall A : |A| > 1$ and get back previous (normal) weighted average, in that

$$\text{WAVG}_{w'}(x) = \text{AG}_w(x) \quad (15.62)$$

Integration, Aggregation, and Weighted Averages

- More complex “nonlinear” aggregation functions can be constructed by defining the aggregation function on all vertices of the hypercube. I.e., for each $\mathbf{1}_A : A \subseteq E$ we might have (for all $A \subseteq E$):

$$\text{AG}(\mathbf{1}_A) = w_A \quad (15.60)$$

- What then might $\text{AG}(x)$ be for some $x \in \mathbb{R}^E$? Our weighted average functions might look **something** more like the r.h.s. in:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.61)$$

- Note, we can define $w(e) = w'(e)$ and $w(A) = 0, \forall A : |A| > 1$ and get back previous (normal) weighted average, in that

$$\text{WAVG}_{w'}(x) = \text{AG}_w(x) \quad (15.62)$$

- Set function $f : 2^E \rightarrow \mathbb{R}$ is a **game** if f is normalized $f(\emptyset) = 0$.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.
- A **Boolean function** f is any function $f : \{0, 1\}^m \rightarrow \{0, 1\}$ and is a **pseudo-Boolean function** if $f : \{0, 1\}^m \rightarrow \mathbb{R}$.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.
- A **Boolean function** f is any function $f : \{0, 1\}^m \rightarrow \{0, 1\}$ and is a **pseudo-Boolean function** if $f : \{0, 1\}^m \rightarrow \mathbb{R}$.
- Any set function corresponds to a pseudo-Boolean function. I.e., given $f : 2^E \rightarrow \mathbb{R}$, form $f_b : \{0, 1\}^m \rightarrow \mathbb{R}$ as $f_b(x) = f(A_x)$ where the A, x bijection is $A = \{e \in E : x_e = 1\}$ and $x = \mathbf{1}_A$.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.
- A **Boolean function** f is any function $f : \{0, 1\}^m \rightarrow \{0, 1\}$ and is a **pseudo-Boolean function** if $f : \{0, 1\}^m \rightarrow \mathbb{R}$.
- Any set function corresponds to a pseudo-Boolean function. I.e., given $f : 2^E \rightarrow \mathbb{R}$, form $f_b : \{0, 1\}^m \rightarrow \mathbb{R}$ as $f_b(x) = f(A_x)$ where the A, x bijection is $A = \{e \in E : x_e = 1\}$ and $x = \mathbf{1}_A$.
- Also, if we have an expression for f_b we can construct a set function f as $f(A) = f_b(\mathbf{1}_A)$. We can also often relax f_b to any $x \in [0, 1]^m$.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.
- A **Boolean function** f is any function $f : \{0, 1\}^m \rightarrow \{0, 1\}$ and is a **pseudo-Boolean function** if $f : \{0, 1\}^m \rightarrow \mathbb{R}$.
- Any set function corresponds to a pseudo-Boolean function. I.e., given $f : 2^E \rightarrow \mathbb{R}$, form $f_b : \{0, 1\}^m \rightarrow \mathbb{R}$ as $f_b(x) = f(A_x)$ where the A, x bijection is $A = \{e \in E : x_e = 1\}$ and $x = \mathbf{1}_A$.
- Also, if we have an expression for f_b we can construct a set function f as $f(A) = f_b(\mathbf{1}_A)$. We can also often relax f_b to any $x \in [0, 1]^m$.
- We saw this for Lovász extension.

Integration, Aggregation, and Weighted Averages

- Set function $f : 2^E \rightarrow \mathbb{R}$ is called a **capacity** if it is monotone non-decreasing, i.e., $f(A) \leq f(B)$ whenever $A \subseteq B$.
- A **Boolean function** f is any function $f : \{0, 1\}^m \rightarrow \{0, 1\}$ and is a **pseudo-Boolean function** if $f : \{0, 1\}^m \rightarrow \mathbb{R}$.
- Any set function corresponds to a pseudo-Boolean function. I.e., given $f : 2^E \rightarrow \mathbb{R}$, form $f_b : \{0, 1\}^m \rightarrow \mathbb{R}$ as $f_b(x) = f(A_x)$ where the A, x bijection is $A = \{e \in E : x_e = 1\}$ and $x = \mathbf{1}_A$.
- Also, if we have an expression for f_b we can construct a set function f as $f(A) = f_b(\mathbf{1}_A)$. We can also often relax f_b to any $x \in [0, 1]^m$.
- We saw this for Lovász extension.
- It turns out that a concept essentially identical to the Lovász extension was derived much earlier, in 1954, and using this derivation (via integration) leads to deeper intuition.

Choquet integral

Definition 15.5.1

Let f be any capacity on E and $w \in \mathbb{R}_+^E$. The **Choquet integral** (1954) of w w.r.t. f is defined by

$$C_f(w) = \sum_{i=1}^m (w_{e_i} - w_{e_{i+1}}) f(E_i) \quad (15.63)$$

where in the sum, we have sorted and renamed the elements of E so that $w_{e_1} \geq w_{e_2} \geq \dots \geq w_{e_m} \geq w_{e_{m+1}} \triangleq 0$, and where $E_i = \{e_1, e_2, \dots, e_i\}$.

- We immediately see that an equivalent formula is as follows:

$$C_f(w) = \sum_{i=1}^m w(e_i) (f(E_i) - f(E_{i-1})) \quad (15.64)$$

where $E_0 \stackrel{\text{def}}{=} \emptyset$.

Choquet integral

Definition 15.5.1

Let f be any capacity on E and $w \in \mathbb{R}_+^E$. The **Choquet integral** (1954) of w w.r.t. f is defined by

$$C_f(w) = \sum_{i=1}^m (w_{e_i} - w_{e_{i+1}}) f(E_i) \quad (15.63)$$

where in the sum, we have sorted and renamed the elements of E so that $w_{e_1} \geq w_{e_2} \geq \dots \geq w_{e_m} \geq w_{e_{m+1}} \triangleq 0$, and where $E_i = \{e_1, e_2, \dots, e_i\}$.

- BTW: this again essentially **Abel's partial summation formula**: Given two arbitrary sequences $\{a_n\}$ and $\{b_n\}$ with $A_n = \sum_{k=1}^n a_k$, we have

$$\sum_{k=m}^n a_k b_k = \sum_{k=m}^n A_k (b_k - b_{k+1}) + A_n b_{n+1} - A_{m-1} b_m \quad (15.65)$$

The “integral” in the Choquet integral

- Thought of as an integral over \mathbb{R} of a piece-wise constant function.

The “integral” in the Choquet integral


- Thought of as an integral over \mathbb{R} of a piece-wise constant function.
- First note, assuming E is ordered according to descending w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_{m-1}) \geq w(e_m)$, then $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e \geq w_{e_i}\}$.

The “integral” in the Choquet integral

- Thought of as an integral over \mathbb{R} of a piece-wise constant function.
- First note, assuming E is ordered according to descending w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_{m-1}) \geq w(e_m)$, then $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e \geq w_{e_i}\}$.
- For any $w_{e_i} > \alpha \geq w_{e_{i+1}}$ we also have $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e > \alpha\}$.


The “integral” in the Choquet integral

- Thought of as an integral over \mathbb{R} of a piece-wise constant function.
- First note, assuming E is ordered according to descending w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_{m-1}) \geq w(e_m)$, then $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e \geq w_{e_i}\}$.
- For any $w_{e_i} > \alpha \geq w_{e_{i+1}}$ we also have $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e > \alpha\}$.
- Consider segmenting the real-axis at boundary points w_{e_i} , right most is w_{e_1} .



The “integral” in the Choquet integral

- Thought of as an integral over \mathbb{R} of a piece-wise constant function.
- First note, assuming E is ordered according to descending w , so that $w(e_1) \geq w(e_2) \geq \dots \geq w(e_{m-1}) \geq w(e_m)$, then $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e \geq w_{e_i}\}$.
- For any $w_{e_i} > \alpha \geq w_{e_{i+1}}$ we also have $E_i = \{e_1, e_2, \dots, e_i\} = \{e \in E : w_e > \alpha\}$.
- Consider segmenting the real-axis at boundary points w_{e_i} , right most is w_{e_1} .



- A function can be defined on a segment of \mathbb{R} , namely $w_{e_i} > \alpha \geq w_{e_{i+1}}$. This function $F_i : [w_{e_{i+1}}, w_{e_i}) \rightarrow \mathbb{R}$ is defined as

$$F_i(\alpha) = f(\{e \in E : w_e > \alpha\}) = f(E_i) \quad (15.66)$$

The “integral” in the Choquet integral

- We can generalize this to multiple segments of \mathbb{R} (for now, take $w \in \mathbb{R}_+^E$). The piecewise-constant function is defined as:


$$F(\alpha) = \begin{cases} f(E) & \text{if } 0 \leq \alpha < w_m \\ f(\{e \in E : w_e > \alpha\}) & \text{if } w_{e_{i+1}} \leq \alpha < w_{e_i}, i \in \{1, \dots, m-1\} \\ 0 (= f(\emptyset)) & \text{if } w_1 < \alpha \end{cases}$$

The “integral” in the Choquet integral

- We can generalize this to multiple segments of \mathbb{R} (for now, take $w \in \mathbb{R}_+^E$). The piecewise-constant function is defined as:

$$F(\alpha) = \begin{cases} f(E) & \text{if } 0 \leq \alpha < w_m \\ f(\{e \in E : w_e > \alpha\}) & \text{if } w_{e_{i+1}} \leq \alpha < w_{e_i}, i \in \{1, \dots, m-1\} \\ 0 (= f(\emptyset)) & \text{if } w_1 < \alpha \end{cases}$$

- Visualizing a piecewise constant function, where the constant values are given by f evaluated on E_i for each i



Note, what is depicted may be a game but not a capacity. Why?

The “integral” in the Choquet integral

- Now consider the integral, with $w \in \mathbb{R}_+^E$, and normalized f so that $f(\emptyset) = 0$. Recall $w_{m+1} \stackrel{\text{def}}{=} 0$.

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.67)$$

The “integral” in the Choquet integral

- Now consider the integral, with $w \in \mathbb{R}_+^E$, and normalized f so that $f(\emptyset) = 0$. Recall $w_{m+1} \stackrel{\text{def}}{=} 0$.

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.67)$$

$$= \int_0^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.68)$$

The “integral” in the Choquet integral

- Now consider the integral, with $w \in \mathbb{R}_+^E$, and normalized f so that $f(\emptyset) = 0$. Recall $w_{m+1} \stackrel{\text{def}}{=} 0$.

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.67)$$

$$= \int_0^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.68)$$

$$= \int_{w_{m+1}}^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.69)$$

The “integral” in the Choquet integral

- Now consider the integral, with $w \in \mathbb{R}_+^E$, and normalized f so that $f(\emptyset) = 0$. Recall $w_{m+1} \stackrel{\text{def}}{=} 0$.

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.67)$$

$$= \int_0^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.68)$$

$$= \int_{w_{m+1}}^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.69)$$

$$= \sum_{i=1}^m \int_{w_{i+1}}^{w_i} f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.70)$$

The “integral” in the Choquet integral

- Now consider the integral, with $w \in \mathbb{R}_+^E$, and normalized f so that $f(\emptyset) = 0$. Recall $w_{m+1} \stackrel{\text{def}}{=} 0$.

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.67)$$

$$= \int_0^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.68)$$

$$= \int_{w_{m+1}}^\infty f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.69)$$

$$= \sum_{i=1}^m \int_{w_{i+1}}^{w_i} f(\{e \in E : w_e > \alpha\}) d\alpha \quad (15.70)$$

$$= \sum_{i=1}^m \int_{w_{i+1}}^{w_i} f(E_i) d\alpha = \sum_{i=1}^m f(E_i)(w_i - w_{i+1}) \quad (15.71)$$

The “integral” in the Choquet integral

- But we saw before that $\sum_{i=1}^m f(E_i)(w_i - w_{i+1})$ is just the Lovász extension of a function f .

The “integral” in the Choquet integral

- But we saw before that $\sum_{i=1}^m f(E_i)(w_i - w_{i+1})$ is just the Lovász extension of a function f .
- Thus, we have the following definition:

Definition 15.5.2

Given $w \in \mathbb{R}_+^E$, the Lovász extension (equivalently Choquet integral) may be defined as follows:

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.72)$$

where the function F is defined as before.

The “integral” in the Choquet integral

- But we saw before that $\sum_{i=1}^m f(E_i)(w_i - w_{i+1})$ is just the Lovász extension of a function f .
- Thus, we have the following definition:

Definition 15.5.2

Given $w \in \mathbb{R}_+^E$, the Lovász extension (equivalently Choquet integral) may be defined as follows:

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.72)$$

where the function F is defined as before.

- Note that it is not necessary in general to require $w \in \mathbb{R}_+^E$ (i.e., we can take $w \in \mathbb{R}^E$) nor that f be non-negative, but it is a bit more involved. Above is the simple case.

The “integral” in the Choquet integral

- But we saw before that $\sum_{i=1}^m f(E_i)(w_i - w_{i+1})$ is just the Lovász extension of a function f .
- Thus, we have the following definition:

Definition 15.5.2

Given $w \in \mathbb{R}_+^E$, the Lovász extension (equivalently Choquet integral) may be defined as follows:

$$\tilde{f}(w) \stackrel{\text{def}}{=} \int_0^\infty F(\alpha) d\alpha \quad (15.72)$$

where the function F is defined as before.

- Note that it is not necessary in general to require $w \in \mathbb{R}_+^E$ (i.e., we can take $w \in \mathbb{R}^E$) nor that f be non-negative, but it is a bit more involved. Above is the simple case.
- The above integral will be further generalized a bit later.

Choquet integral and aggregation

- Recall, we want to produce some notion of generalized aggregation function having the flavor of:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.73)$$

how does this correspond to Lovász extension?

Choquet integral and aggregation

- Recall, we want to produce some notion of generalized aggregation function having the flavor of:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.73)$$

how does this correspond to Lovász extension?

- Let us partition the hypercube $[0, 1]^m$ into q polytopes, each defined by a set of vertices $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_q$.

Choquet integral and aggregation

- Recall, we want to produce some notion of generalized aggregation function having the flavor of:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.73)$$

how does this correspond to Lovász extension?

- Let us partition the hypercube $[0, 1]^m$ into q polytopes, each defined by a set of vertices $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_q$.
- E.g., for each i , $\mathcal{V}_i = \{\mathbf{1}_{A_1}, \mathbf{1}_{A_2}, \dots, \mathbf{1}_{A_k}\}$ (k vertices) and the convex hull of \mathcal{V}_i defines the i^{th} polytope.

Choquet integral and aggregation

- Recall, we want to produce some notion of generalized aggregation function having the flavor of:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.73)$$

how does this correspond to Lovász extension?

- Let us partition the hypercube $[0, 1]^m$ into q polytopes, each defined by a set of vertices $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_q$.
- E.g., for each i , $\mathcal{V}_i = \{\mathbf{1}_{A_1}, \mathbf{1}_{A_2}, \dots, \mathbf{1}_{A_k}\}$ (k vertices) and the convex hull of \mathcal{V}_i defines the i^{th} polytope.
- This forms a “triangulation” of the hypercube.

Choquet integral and aggregation

- Recall, we want to produce some notion of generalized aggregation function having the flavor of:

$$\text{AG}(x) = \sum_{A \subseteq E} x(A)w_A = \sum_{A \subseteq E} x(A)\text{AG}(\mathbf{1}_A) \quad (15.73)$$

how does this correspond to Lovász extension?

- Let us partition the hypercube $[0, 1]^m$ into q polytopes, each defined by a set of vertices $\mathcal{V}_1, \mathcal{V}_2, \dots, \mathcal{V}_q$.
- E.g., for each i , $\mathcal{V}_i = \{\mathbf{1}_{A_1}, \mathbf{1}_{A_2}, \dots, \mathbf{1}_{A_k}\}$ (k vertices) and the convex hull of \mathcal{V}_i defines the i^{th} polytope.
- This forms a “triangulation” of the hypercube.
- For any $x \in [0, 1]^m$ there is a (not necessarily unique) $\mathcal{V}(x) = \mathcal{V}_j$ for some j such that $x \in \text{conv}(\mathcal{V}(x))$.

Choquet integral and aggregation

- Most generally, for $x \in [0, 1]^m$, let us define the (unique) coefficients $\alpha_0^x(A)$ and $\alpha_i^x(A)$ that define the affine transformation of the coefficients of x to be used with the particular hypercube vertex $\mathbf{1}_A \in \text{conv}(\mathcal{V}(x))$. The affine transformation is as follows:

$$\alpha_0^x(A) + \sum_{j=1}^m \alpha_j^x(A)x_j \in \mathbb{R} \quad (15.74)$$

Note that many of these coefficient are often zero.

Choquet integral and aggregation

- Most generally, for $x \in [0, 1]^m$, let us define the (unique) coefficients $\alpha_0^x(A)$ and $\alpha_i^x(A)$ that define the affine transformation of the coefficients of x to be used with the particular hypercube vertex $\mathbf{1}_A \in \text{conv}(\mathcal{V}(x))$. The affine transformation is as follows:

$$\alpha_0^x(A) + \sum_{j=1}^m \alpha_j^x(A)x_j \in \mathbb{R} \quad (15.74)$$

Note that many of these coefficient are often zero.

- From this, we can define an aggregation function of the form

$$\text{AG}(x) \stackrel{\text{def}}{=} \sum_{A: \mathbf{1}_A \in \mathcal{V}(x)} \left(\alpha_0^x(A) + \sum_{j=1}^m \alpha_j^x(A)x_j \right) \text{AG}(\mathbf{1}_A) \quad (15.75)$$

Choquet integral and aggregation

- We can define a canonical triangulation of the hypercube in terms of permutations of the coordinates. I.e., given some permutation σ , define

$$\text{conv}(\mathcal{V}_\sigma) = \{x \in [0, 1]^n \mid x_{\sigma(1)} \geq x_{\sigma(2)} \geq \cdots \geq x_{\sigma(m)}\} \quad (15.76)$$

Then these $m!$ blocks of the partition are called the **canonical partitions** of the hypercube.

Choquet integral and aggregation

- We can define a canonical triangulation of the hypercube in terms of permutations of the coordinates. I.e., given some permutation σ , define

$$\text{conv}(\mathcal{V}_\sigma) = \{x \in [0, 1]^n \mid x_{\sigma(1)} \geq x_{\sigma(2)} \geq \cdots \geq x_{\sigma(m)}\} \quad (15.76)$$

Then these $m!$ blocks of the partition are called the **canonical partitions** of the hypercube.

- With this, we can define $\{\mathcal{V}_i\}_i$ as the vertices of $\text{conv}(\mathcal{V}_\sigma)$ for each permutation σ .

Choquet integral and aggregation

- We can define a canonical triangulation of the hypercube in terms of permutations of the coordinates. I.e., given some permutation σ , define

$$\text{conv}(\mathcal{V}_\sigma) = \{x \in [0, 1]^n \mid x_{\sigma(1)} \geq x_{\sigma(2)} \geq \cdots \geq x_{\sigma(m)}\} \quad (15.76)$$

Then these $m!$ blocks of the partition are called the **canonical partitions** of the hypercube.

- With this, we can define $\{\mathcal{V}_i\}_i$ as the vertices of $\text{conv}(\mathcal{V}_\sigma)$ for each permutation σ . **In this case, we have:**

Choquet integral and aggregation

- We can define a canonical triangulation of the hypercube in terms of permutations of the coordinates. I.e., given some permutation σ , define

$$\text{conv}(\mathcal{V}_\sigma) = \{x \in [0, 1]^n \mid x_{\sigma(1)} \geq x_{\sigma(2)} \geq \cdots \geq x_{\sigma(m)}\} \quad (15.76)$$

Then these $m!$ blocks of the partition are called the **canonical partitions** of the hypercube.

- With this, we can define $\{\mathcal{V}_i\}_i$ as the vertices of $\text{conv}(\mathcal{V}_\sigma)$ for each permutation σ . In this case, we have:

Proposition 15.5.3

The above linear interpolation in Eqn. (15.75) using the canonical partition yields the Lovász extension with $\alpha_0^x(A) + \sum_{j=1}^m \alpha_j^x(A)x_j = x_{\sigma_i} - x_{\sigma_{i-1}}$ for $A = E_i = \{e_{\sigma_1}, \dots, e_{\sigma_i}\}$ for appropriate order σ .

Choquet integral and aggregation

- We can define a canonical triangulation of the hypercube in terms of permutations of the coordinates. I.e., given some permutation σ , define

$$\text{conv}(\mathcal{V}_\sigma) = \{x \in [0, 1]^n \mid x_{\sigma(1)} \geq x_{\sigma(2)} \geq \cdots \geq x_{\sigma(m)}\} \quad (15.76)$$

Then these $m!$ blocks of the partition are called the **canonical partitions** of the hypercube.

- With this, we can define $\{\mathcal{V}_i\}_i$ as the vertices of $\text{conv}(\mathcal{V}_\sigma)$ for each permutation σ . In this case, we have:

Proposition 15.5.3

The above linear interpolation in Eqn. (15.75) using the canonical partition yields the Lovász extension with $\alpha_0^x(A) + \sum_{j=1}^m \alpha_j^x(A)x_j = x_{\sigma_i} - x_{\sigma_{i-1}}$ for $A = E_i = \{e_{\sigma_1}, \dots, e_{\sigma_i}\}$ for appropriate order σ .

- Hence, Lovász extension is a generalized aggregation function.

Lovász extension as max over orders

- We can also write the Lovász extension as follows:

$$\tilde{f}(w) = \max_{\sigma \in \Pi_{[m]}} w^\top c^\sigma \quad (15.77)$$

where $\Pi_{[m]}$ is the set of $m!$ permutations of $[m] = E$, $\sigma \in \Pi_{[m]}$ is a particular permutation, and c^σ is a vector associated with permutation σ defined as:

$$c_i^\sigma = f(E_{\sigma_i}) - f(E_{\sigma_{i-1}}) \quad (15.78)$$

where $E_{\sigma_i} = \{e_{\sigma_1}, e_{\sigma_2}, \dots, e_{\sigma_i}\}$.

Lovász extension as max over orders

- We can also write the Lovász extension as follows:

$$\tilde{f}(w) = \max_{\sigma \in \Pi_{[m]}} w^\top c^\sigma \quad (15.77)$$

where $\Pi_{[m]}$ is the set of $m!$ permutations of $[m] = E$, $\sigma \in \Pi_{[m]}$ is a particular permutation, and c^σ is a vector associated with permutation σ defined as:

$$c_i^\sigma = f(E_{\sigma_i}) - f(E_{\sigma_{i-1}}) \quad (15.78)$$

where $E_{\sigma_i} = \{e_{\sigma_1}, e_{\sigma_2}, \dots, e_{\sigma_i}\}$.

- Note this immediately follows from the definition of the Lovász extension in the form:

$$\tilde{f}(w) = \max_{x \in P_f} w^\top x = \max_{x \in B_f} w^\top x \quad (15.79)$$

since we know that the maximum is achieved by an extreme point of the base B_f and all extreme points are obtained by a permutation-of- E -parameterized greedy instance.

Lovász extension, defined in multiple ways

- As shorthand notation, lets use $\{w \geq \alpha\} \equiv \{e \in E : w(e) \geq \alpha\}$, called the weak α -sup-level set of w .

Lovász extension, defined in multiple ways

- As shorthand notation, lets use $\{w \geq \alpha\} \equiv \{e \in E : w(e) \geq \alpha\}$, called the weak α -sup-level set of w . A similar definition holds for $\{w > \alpha\}$ (called the strong α -sup-level set of w).

Lovász extension, defined in multiple ways

- As shorthand notation, lets use $\{w \geq \alpha\} \equiv \{e \in E : w(e) \geq \alpha\}$, called the weak α -sup-level set of w . A similar definition holds for $\{w > \alpha\}$ (called the strong α -sup-level set of w).
- Given **any** $w \in \mathbb{R}^E$, sort E as $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$.

Lovász extension, defined in multiple ways

- As shorthand notation, lets use $\{w \geq \alpha\} \equiv \{e \in E : w(e) \geq \alpha\}$, called the weak α -sup-level set of w . A similar definition holds for $\{w > \alpha\}$ (called the strong α -sup-level set of w).
- Given **any** $w \in \mathbb{R}^E$, sort E as $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$. **Also**, w.l.o.g., number elements of w so that $w_1 \geq w_2 \geq \dots \geq w_m$.

Lovász extension, defined in multiple ways

- As shorthand notation, let's use $\{w \geq \alpha\} \equiv \{e \in E : w(e) \geq \alpha\}$, called the weak α -sup-level set of w . A similar definition holds for $\{w > \alpha\}$ (called the strong α -sup-level set of w).
- Given **any** $w \in \mathbb{R}^E$, sort E as $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m)$. Also, w.l.o.g., number elements of w so that $w_1 \geq w_2 \geq \dots \geq w_m$.
- We have already seen how we can define the Lovász extension for any (not necessarily submodular) function f in the following equivalent ways:

$$\tilde{f}(w) = \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) \quad (15.80)$$

$$= \sum_{i=1}^{m-1} f(E_i)(w(e_i) - w(e_{i+1})) + f(E)w(e_m)a \quad (15.81)$$

$$= \sum_{i=1}^{m-1} \lambda_i f(E_i) \quad (15.82)$$

Lovász extension, as integral

- Additional ways we can define the Lovász extension for any (not necessarily submodular) but normalized function f include:

$$\tilde{f}(w) = \sum_{i=1}^m w(e_i) f(e_i | E_{i-1}) = \sum_{i=1}^m \lambda_i f(E_i) \quad (15.83)$$

$$= \sum_{i=1}^{m-1} f(E_i)(w(e_i) - w(e_{i+1})) + f(E)w(e_m) \quad (15.84)$$

$$= \int_{\min\{w_1, \dots, w_m\}}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min\{w_1, \dots, w_m\} \quad (15.85)$$

$$= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{-\infty}^0 [f(\{w \geq \alpha\}) - f(E)] d\alpha \quad (15.86)$$

general Lovász extension, as simple integral

- In fact, we have that, given function f , and any $w \in \mathbb{R}^E$:

$$\tilde{f}(w) = \int_{-\infty}^{+\infty} \hat{f}(\alpha) d\alpha \quad (15.87)$$

where

$$\hat{f}(\alpha) = \begin{cases} f(\{w \geq \alpha\}) & \text{if } \alpha \geq 0 \\ f(\{w \geq \alpha\}) - f(E) & \text{if } \alpha < 0 \end{cases} \quad (15.88)$$

general Lovász extension, as simple integral

- In fact, we have that, given function f , and any $w \in \mathbb{R}^E$:

$$\tilde{f}(w) = \int_{-\infty}^{+\infty} \hat{f}(\alpha) d\alpha \quad (15.87)$$

where

$$\hat{f}(\alpha) = \begin{cases} f(\{w \geq \alpha\}) & \text{if } \alpha \geq 0 \\ f(\{w \geq \alpha\}) - f(E) & \text{if } \alpha < 0 \end{cases} \quad (15.88)$$

- So we can write it as a simple integral over the right function.

general Lovász extension, as simple integral

- In fact, we have that, given function f , and any $w \in \mathbb{R}^E$:

$$\tilde{f}(w) = \int_{-\infty}^{+\infty} \hat{f}(\alpha) d\alpha \quad (15.87)$$

where

$$\hat{f}(\alpha) = \begin{cases} f(\{w \geq \alpha\}) & \text{if } \alpha \geq 0 \\ f(\{w \geq \alpha\}) - f(E) & \text{if } \alpha < 0 \end{cases} \quad (15.88)$$

- So we can write it as a simple integral over the right function.
- These make it easier to see certain properties of the Lovász extension. But first, we show the above.

Lovász extension, as integral

- To show Eqn. (15.85), first note that the r.h.s. terms are the same since $w(e_m) = \min \{w_1, \dots, w_m\}$.

Lovász extension, as integral

- To show Eqn. (15.85), first note that the r.h.s. terms are the same since $w(e_m) = \min \{w_1, \dots, w_m\}$.
- Then, consider that, as a function of α , we have

$$f(\{w \geq \alpha\}) = \begin{cases} 0 & \text{if } \alpha > w(e_1) \\ f(E_k) & \text{if } \alpha \in (w(e_{k+1}), w(e_k)), k \in \{1, \dots, m-1\} \\ f(E) & \text{if } \alpha < w(e_m) \end{cases} \quad (15.89)$$

we may use open intervals since sets of zero measure don't change integration.

Lovász extension, as integral

- To show Eqn. (15.85), first note that the r.h.s. terms are the same since $w(e_m) = \min \{w_1, \dots, w_m\}$.
- Then, consider that, as a function of α , we have

$$f(\{w \geq \alpha\}) = \begin{cases} 0 & \text{if } \alpha > w(e_1) \\ f(E_k) & \text{if } \alpha \in (w(e_{k+1}), w(e_k)), k \in \{1, \dots, m-1\} \\ f(E) & \text{if } \alpha < w(e_m) \end{cases} \quad (15.89)$$

we may use open intervals since sets of zero measure don't change integration.

- Inside the integral, then, this recovers Eqn. (15.84).

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\tilde{f}(w)$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\tilde{f}(w) = \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\}$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned} \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\ &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \end{aligned}$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned} \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\ &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \\ &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(E) d\alpha + \int_0^{w_m} f(E) d\alpha \end{aligned}$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned}
 \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(E) d\alpha + \int_0^{w_m} f(E) d\alpha \\
 &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^0 f(E) d\alpha
 \end{aligned}$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned}
 \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(E) d\alpha + \int_0^{w_m} f(E) d\alpha \\
 &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^0 f(E) d\alpha \\
 &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 [f(\{w \geq \alpha\}) - f(E)] d\alpha
 \end{aligned}$$

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned}
 \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \\
 &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(E) d\alpha + \int_0^{w_m} f(E) d\alpha \\
 &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^0 f(E) d\alpha \\
 &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 [f(\{w \geq \alpha\}) - f(E)] d\alpha
 \end{aligned}$$

and then let $\beta \rightarrow \infty$ and we get Eqn. (15.86), i.e.:

Lovász extension, as integral

- To show Eqn. (15.86), start with Eqn. (15.85), note

$w_m = \min \{w_1, \dots, w_m\}$, take any $\beta \leq \min \{0, w_1, \dots, w_m\}$, and form:

$$\begin{aligned} \tilde{f}(w) &= \int_{w_m}^{+\infty} f(\{w \geq \alpha\}) d\alpha + f(E) \min \{w_1, \dots, w_m\} \\ &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(\{w \geq \alpha\}) d\alpha + f(E) \int_0^{w_m} d\alpha \\ &= \int_{\beta}^{+\infty} f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^{w_m} f(E) d\alpha + \int_0^{w_m} f(E) d\alpha \\ &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 f(\{w \geq \alpha\}) d\alpha - \int_{\beta}^0 f(E) d\alpha \\ &= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{\beta}^0 [f(\{w \geq \alpha\}) - f(E)] d\alpha \end{aligned}$$

and then let $\beta \rightarrow \infty$ and we get Eqn. (15.86), i.e.:

$$= \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha + \int_{-\infty}^0 [f(\{w \geq \alpha\}) - f(E)] d\alpha$$

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1 *Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.*

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.
- 2** If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\})d\alpha$.

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** *Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.*
- 2** *If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\})d\alpha$.*
- 3** *For $w \in \mathbb{R}^E$, and $\alpha \in \mathbb{R}$, we have $\tilde{f}(w + \alpha\mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.*

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.
- 2** If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha$.
- 3** For $w \in \mathbb{R}^E$, and $\alpha \in \mathbb{R}$, we have $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- 4** Positive homogeneity: I.e., $\tilde{f}(\alpha w) = \alpha \tilde{f}(w)$ for $\alpha \geq 0$.

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.
- 2** If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha$.
- 3** For $w \in \mathbb{R}^E$, and $\alpha \in \mathbb{R}$, we have $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- 4** Positive homogeneity: I.e., $\tilde{f}(\alpha w) = \alpha \tilde{f}(w)$ for $\alpha \geq 0$.
- 5** For all $A \subseteq E$, $\tilde{f}(\mathbf{1}_A) = f(A)$.

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.
- 2** If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha$.
- 3** For $w \in \mathbb{R}^E$, and $\alpha \in \mathbb{R}$, we have $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- 4** Positive homogeneity: I.e., $\tilde{f}(\alpha w) = \alpha \tilde{f}(w)$ for $\alpha \geq 0$.
- 5** For all $A \subseteq E$, $\tilde{f}(\mathbf{1}_A) = f(A)$.
- 6** f symmetric as in $f(A) = f(E \setminus A), \forall A$, then $\tilde{f}(w) = \tilde{f}(-w)$ (\tilde{f} is even).

Lovász extension properties

- Using the above, have the following (some of which we've seen):

Theorem 15.6.1

Let $f, g : 2^E \rightarrow \mathbb{R}$ be normalized ($f(\emptyset) = g(\emptyset) = 0$). Then

- 1** Superposition of LE operator: Given f and g with Lovász extensions \tilde{f} and \tilde{g} then $\tilde{f} + \tilde{g}$ is the Lovász extension of $f + g$ and $\lambda\tilde{f}$ is the Lovász extension of λf for $\lambda \in \mathbb{R}$.
- 2** If $w \in \mathbb{R}_+^E$ then $\tilde{f}(w) = \int_0^{+\infty} f(\{w \geq \alpha\}) d\alpha$.
- 3** For $w \in \mathbb{R}^E$, and $\alpha \in \mathbb{R}$, we have $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- 4** Positive homogeneity: I.e., $\tilde{f}(\alpha w) = \alpha \tilde{f}(w)$ for $\alpha \geq 0$.
- 5** For all $A \subseteq E$, $\tilde{f}(\mathbf{1}_A) = f(A)$.
- 6** f symmetric as in $f(A) = f(E \setminus A), \forall A$, then $\tilde{f}(w) = \tilde{f}(-w)$ (\tilde{f} is even).
- 7** Given partition $E^1 \cup E^2 \cup \dots \cup E^k$ of E and $w = \sum_{i=1}^k \gamma_i \mathbf{1}_{E^i}$ with $\gamma_1 \geq \gamma_2 \geq \dots \geq \gamma_k$, and with $E^{1:i} = E^1 \cup E^2 \cup \dots \cup E^i$, then
$$\tilde{f}(w) = \sum_{i=1}^k \gamma_i f(E^i | E^{1:i-1}) = \sum_{i=1}^{k-1} f(E^{1:i})(\gamma_i - \gamma_{i+1}) + f(E)\gamma_k.$$

Lovász extension properties: ex. property 3

- Consider property property 3, for example, which says that $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.

Lovász extension properties: ex. property 3

- Consider property property 3, for example, which says that $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- This means that, say when $m = 2$, that as we move along the line $w_1 = w_2$, the Lovász extension scales linearly.

Lovász extension properties: ex. property 3

- Consider property property 3, for example, which says that $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$.
- This means that, say when $m = 2$, that as we move along the line $w_1 = w_2$, the Lovász extension scales linearly.
- And if $f(E) = 0$, then the Lovász extension is constant along the direction $\mathbf{1}_E$.

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w)$$

(15.92)

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\}) d\alpha$$

(15.92)

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

(15.92)

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

$$\stackrel{(a)}{=} \int_{-\infty}^{\infty} f(\{w \leq \alpha\})d\alpha$$

(15.92)

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

$$\stackrel{(a)}{=} \int_{-\infty}^{\infty} f(\{w \leq \alpha\})d\alpha \stackrel{(b)}{=} \int_{-\infty}^{\infty} f(\{w > \alpha\})d\alpha \quad (15.91)$$

$$(15.92)$$

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

$$\stackrel{(a)}{=} \int_{-\infty}^{\infty} f(\{w \leq \alpha\})d\alpha \stackrel{(b)}{=} \int_{-\infty}^{\infty} f(\{w > \alpha\})d\alpha \quad (15.91)$$

$$= \int_{-\infty}^{\infty} f(\{w \geq \alpha\})d\alpha \quad (15.92)$$

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

$$\stackrel{(a)}{=} \int_{-\infty}^{\infty} f(\{w \leq \alpha\})d\alpha \stackrel{(b)}{=} \int_{-\infty}^{\infty} f(\{w > \alpha\})d\alpha \quad (15.91)$$

$$= \int_{-\infty}^{\infty} f(\{w \geq \alpha\})d\alpha = \tilde{f}(w) \quad (15.92)$$

Lovász extension properties

- Given Eqns. (15.83) through (15.86), most of the above properties are relatively easy to derive.
- For example, if f is symmetric, and since $f(E) = f(\emptyset) = 0$, we have

$$\tilde{f}(-w) = \int_{-\infty}^{\infty} f(\{-w \geq \alpha\})d\alpha = \int_{-\infty}^{\infty} f(\{w \leq -\alpha\})d\alpha \quad (15.90)$$

$$\stackrel{(a)}{=} \int_{-\infty}^{\infty} f(\{w \leq \alpha\})d\alpha \stackrel{(b)}{=} \int_{-\infty}^{\infty} f(\{w > \alpha\})d\alpha \quad (15.91)$$

$$= \int_{-\infty}^{\infty} f(\{w \geq \alpha\})d\alpha = \tilde{f}(w) \quad (15.92)$$

Equality (a) follows since $\int_{-\infty}^{\infty} f(\alpha)d\alpha = \int_{-\infty}^{\infty} f(a\alpha + b)d\alpha$ for any b and $a \in \pm 1$, and equality (b) follows since $f(A) = f(E \setminus A)$, so $f(\{w \leq \alpha\}) = f(\{w > \alpha\})$.

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$
- Since $f(\{w \geq \alpha\}) = 0$ for $\alpha > w_1 \geq w_\ell$, we have for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha$

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$
- Since $f(\{w \geq \alpha\}) = 0$ for $\alpha > w_1 \geq w_\ell$, we have for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha$
- For $w \in [0, 1]^E$, then $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha = \int_0^1 f(\{w \geq \alpha\})d\alpha$ since $f(\{w \geq \alpha\}) = 0$ for $1 \geq \alpha > w_1$.

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$
- Since $f(\{w \geq \alpha\}) = 0$ for $\alpha > w_1 \geq w_\ell$, we have for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha$
- For $w \in [0, 1]^E$, then $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha = \int_0^1 f(\{w \geq \alpha\})d\alpha$ since $f(\{w \geq \alpha\}) = 0$ for $1 \geq \alpha > w_1$.
- Consider α as a uniform random variable on $[0, 1]$ and let $h(\alpha)$ be a function of α . Then the expected value $\mathbb{E}[h(\alpha)] = \int_0^1 h(\alpha)d\alpha$.

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$
- Since $f(\{w \geq \alpha\}) = 0$ for $\alpha > w_1 \geq w_\ell$, we have for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha$
- For $w \in [0, 1]^E$, then $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha = \int_0^1 f(\{w \geq \alpha\})d\alpha$ since $f(\{w \geq \alpha\}) = 0$ for $1 \geq \alpha > w_1$.
- Consider α as a uniform random variable on $[0, 1]$ and let $h(\alpha)$ be a function of α . Then the expected value $\mathbb{E}[h(\alpha)] = \int_0^1 h(\alpha)d\alpha$.
- Hence, for $w \in [0, 1]^m$, we can also define the Lovász extension as

$$\tilde{f}(w) = \mathbb{E}_{p(\alpha)}[\underbrace{f(\{w \geq \alpha\})}_{h(\alpha)}] = \mathbb{E}_{p(\alpha)}[\underbrace{f(e \in E : w(e_i) \geq \alpha)}_{h(\alpha)}] \quad (15.93)$$

where α is uniform random variable in $[0, 1]$.

Lovász extension, expected value of random variable

- Recall, for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^\infty f(\{w \geq \alpha\})d\alpha$
- Since $f(\{w \geq \alpha\}) = 0$ for $\alpha > w_1 \geq w_\ell$, we have for $w \in \mathbb{R}_+^E$, we have $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha$
- For $w \in [0, 1]^E$, then $\tilde{f}(w) = \int_0^{w_1} f(\{w \geq \alpha\})d\alpha = \int_0^1 f(\{w \geq \alpha\})d\alpha$ since $f(\{w \geq \alpha\}) = 0$ for $1 \geq \alpha > w_1$.
- Consider α as a uniform random variable on $[0, 1]$ and let $h(\alpha)$ be a function of α . Then the expected value $\mathbb{E}[h(\alpha)] = \int_0^1 h(\alpha)d\alpha$.
- Hence, for $w \in [0, 1]^m$, we can also define the Lovász extension as

$$\tilde{f}(w) = \mathbb{E}_{p(\alpha)}[\underbrace{f(\{w \geq \alpha\})}_{h(\alpha)}] = \mathbb{E}_{p(\alpha)}[\underbrace{f(e \in E : w(e_i) \geq \alpha)}_{h(\alpha)}] \quad (15.93)$$

where α is uniform random variable in $[0, 1]$.

- Useful for showing results for randomized rounding schemes in solving submodular opt. problems subject to constraints via relaxations to convex optimization problems subject to linear constraints.

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.94)$$

$$= (w_1 - w_2) f(\{1\}) + w_2 f(\{1, 2\}) \quad (15.95)$$

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.94)$$

$$= (w_1 - w_2) f(\{1\}) + w_2 f(\{1, 2\}) \quad (15.95)$$

- If $w_1 \leq w_2$, then

$$\tilde{f}(w) = w_2 f(\{2\}) + w_1 f(\{1\}|\{2\}) \quad (15.96)$$

$$= (w_2 - w_1) f(\{2\}) + w_1 f(\{1, 2\}) \quad (15.97)$$

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.98)$$

$$= (w_1 - w_2) f(\{1\}) + w_2 f(\{1, 2\}) \quad (15.99)$$

$$= \frac{1}{2} f(1)(w_1 - w_2) + \frac{1}{2} f(1)(w_1 - w_2) \quad (15.100)$$

$$+ \frac{1}{2} f(\{1, 2\})(w_1 + w_2) - \frac{1}{2} f(\{1, 2\})(w_1 - w_2) \quad (15.101)$$

$$+ \frac{1}{2} f(2)(w_1 - w_2) + \frac{1}{2} f(2)(w_2 - w_1) \quad (15.102)$$

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.98)$$

$$= (w_1 - w_2) f(\{1\}) + w_2 f(\{1, 2\}) \quad (15.99)$$

$$= \frac{1}{2} f(1)(w_1 - w_2) + \frac{1}{2} f(1)(w_1 - w_2) \quad (15.100)$$

$$+ \frac{1}{2} f(\{1, 2\})(w_1 + w_2) - \frac{1}{2} f(\{1, 2\})(w_1 - w_2) \quad (15.101)$$

$$+ \frac{1}{2} f(2)(w_1 - w_2) + \frac{1}{2} f(2)(w_2 - w_1) \quad (15.102)$$

- A similar (symmetric) expression holds when $w_1 \leq w_2$.

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- This gives, for general w_1, w_2 , that

$$\tilde{f}(w) = \frac{1}{2} (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) |w_1 - w_2| \quad (15.103)$$

$$+ \frac{1}{2} (f(\{1\}) - f(\{2\}) + f(\{1, 2\})) w_1 \quad (15.104)$$

$$+ \frac{1}{2} (-f(\{1\}) + f(\{2\}) + f(\{1, 2\})) w_2 \quad (15.105)$$

$$= - (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) \min \{w_1, w_2\} \quad (15.106)$$

$$+ f(\{1\})w_1 + f(\{2\})w_2 \quad (15.107)$$

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- This gives, for general w_1, w_2 , that

$$\tilde{f}(w) = \frac{1}{2} (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) |w_1 - w_2| \quad (15.103)$$

$$+ \frac{1}{2} (f(\{1\}) - f(\{2\}) + f(\{1, 2\})) w_1 \quad (15.104)$$

$$+ \frac{1}{2} (-f(\{1\}) + f(\{2\}) + f(\{1, 2\})) w_2 \quad (15.105)$$

$$= - (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) \min \{w_1, w_2\} \quad (15.106)$$

$$+ f(\{1\})w_1 + f(\{2\})w_2 \quad (15.107)$$

- Thus, if $f(A) = H(X_A)$ is the entropy function, we have

$\tilde{f}(w) = H(e_1)w_1 + H(e_2)w_2 - I(e_1; e_2) \min \{w_1, w_2\}$ which must be convex in w , where $I(e_1; e_2)$ is the mutual information.

Simple expressions for Lovász E . with $m = 2$, $E = \{1, 2\}$

- This gives, for general w_1, w_2 , that

$$\tilde{f}(w) = \frac{1}{2} (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) |w_1 - w_2| \quad (15.103)$$

$$+ \frac{1}{2} (f(\{1\}) - f(\{2\}) + f(\{1, 2\})) w_1 \quad (15.104)$$

$$+ \frac{1}{2} (-f(\{1\}) + f(\{2\}) + f(\{1, 2\})) w_2 \quad (15.105)$$

$$= - (f(\{1\}) + f(\{2\}) - f(\{1, 2\})) \min \{w_1, w_2\} \quad (15.106)$$

$$+ f(\{1\})w_1 + f(\{2\})w_2 \quad (15.107)$$

- Thus, if $f(A) = H(X_A)$ is the entropy function, we have $\tilde{f}(w) = H(e_1)w_1 + H(e_2)w_2 - I(e_1; e_2) \min \{w_1, w_2\}$ which must be convex in w , where $I(e_1; e_2)$ is the mutual information.
- This “simple” but general form of the Lovász extension with $m = 2$ can be useful.

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

- If $w_1 \leq w_2$, then

$$\tilde{f}(w) = w_2 f(\{2\}) + w_1 f(\{1\}|\{2\}) \quad (15.109)$$

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

- If $w_1 \leq w_2$, then

$$\tilde{f}(w) = w_2 f(\{2\}) + w_1 f(\{1\}|\{2\}) \quad (15.109)$$

- If $w = (0, 1)/f(\{2\}) = (0, 1/f(\{2\}))$ then $\tilde{f}(w) = 1$.

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

- If $w_1 \leq w_2$, then

$$\tilde{f}(w) = w_2 f(\{2\}) + w_1 f(\{1\}|\{2\}) \quad (15.109)$$

- If $w = (0, 1)/f(\{2\}) = (0, 1/f(\{2\}))$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

Example: $m = 2$, $E = \{1, 2\}$, contours

- If $w_1 \geq w_2$, then

$$\tilde{f}(w) = w_1 f(\{1\}) + w_2 f(\{2\}|\{1\}) \quad (15.108)$$

- If $w = (1, 0)/f(\{1\}) = (1/f(\{1\}), 0)$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

- If $w_1 \leq w_2$, then


$$\tilde{f}(w) = w_2 f(\{2\}) + w_1 f(\{1\}|\{2\}) \quad (15.109)$$

- If $w = (0, 1)/f(\{2\}) = (0, 1/f(\{2\}))$ then $\tilde{f}(w) = 1$.
- If $w = (1, 1)/f(\{1, 2\})$ then $\tilde{f}(w) = 1$.

- Can plot contours of the form $\left\{ w \in \mathbb{R}^2 : \tilde{f}(w) = 1 \right\}$, particular marked points of form $w = \mathbf{1}_A \times \frac{1}{f(A)}$ for certain A , where $\tilde{f}(w) = 1$.

Example: $m = 2$, $E = \{1, 2\}$

- Contour plot of $m = 2$ Lovász extension (from Bach-2011).



Example: $m = 3$, $E = \{1, 2, 3\}$

- In order to visualize in 3D, we make a few simplifications.

Example: $m = 3$, $E = \{1, 2, 3\}$

- In order to visualize in 3D, we make a few simplifications.
- Consider any submodular f' and $x \in B_{f'}$. Then $f(A) = f'(A) - x(A)$ is submodular

Example: $m = 3$, $E = \{1, 2, 3\}$

- In order to visualize in 3D, we make a few simplifications.
- Consider any submodular f' and $x \in B_{f'}$. Then $f(A) = f'(A) - x(A)$ is submodular, and moreover $f(E) = f'(E) - x(E) = 0$.

Example: $m = 3$, $E = \{1, 2, 3\}$

- In order to visualize in 3D, we make a few simplifications.
- Consider any submodular f' and $x \in B_{f'}$. Then $f(A) = f'(A) - x(A)$ is submodular, and moreover $f(E) = f'(E) - x(E) = 0$.
- Hence, from $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$, we have that $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w)$.

Example: $m = 3$, $E = \{1, 2, 3\}$


- In order to visualize in 3D, we make a few simplifications.
- Consider any submodular f' and $x \in B_{f'}$. Then $f(A) = f'(A) - x(A)$ is submodular, and moreover $f(E) = f'(E) - x(E) = 0$.
- Hence, from $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w) + \alpha f(E)$, we have that $\tilde{f}(w + \alpha \mathbf{1}_E) = \tilde{f}(w)$.
- Thus, we can look “down” on the contour plot of the Lovász extension, $\{w : \tilde{f}(w) = 1\}$, from a vantage point right on the line $\{x : x = \alpha \mathbf{1}_E, \alpha > 0\}$ since moving in direction $\mathbf{1}_E$ changes nothing.

Example: $m = 3$, $E = \{1, 2, 3\}$

- Example 1 (from Bach-2011): $f(A) = \mathbf{1}_{|A| \in \{1,2\}}$
 $= \min\{|A|, 1\} + \min\{|E \setminus A|, 1\} - 1$ is submodular, and
 $\tilde{f}(w) = \max_{k \in \{1,2,3\}} w_k - \min_{k \in \{1,2,3\}} w_k$.


Example: $m = 3$, $E = \{1, 2, 3\}$

- Example 1 (from Bach-2011): $f(A) = \mathbf{1}_{|A| \in \{1,2\}}$
 $= \min\{|A|, 1\} + \min\{|E \setminus A|, 1\} - 1$ is submodular, and
 $\tilde{f}(w) = \max_{k \in \{1,2,3\}} w_k - \min_{k \in \{1,2,3\}} w_k$.




Example: $m = 3$, $E = \{1, 2, 3\}$

- Example 2 (from Bach-2011): $f(A) = |\mathbf{1}_{1 \in A} - \mathbf{1}_{2 \in A}| + |\mathbf{1}_{2 \in A} - \mathbf{1}_{3 \in A}|$







Example: $m = 3$, $E = \{1, 2, 3\}$

- Example 2 (from Bach-2011): $f(A) = |\mathbf{1}_{1 \in A} - \mathbf{1}_{2 \in A}| + |\mathbf{1}_{2 \in A} - \mathbf{1}_{3 \in A}|$
- This gives a “total variation” function for the Lovász extension, with $\tilde{f}(w) = |w_1 - w_2| + |w_2 - w_3|$, a prior to prefer piecewise-constant signals.



Total Variation Example

From “Nonlinear total variation based noise removal algorithms”
Rudin, Osher, and Fatemi, 1992. Top left original, bottom right total variation.



Example: Lovász extension of concave over modular

- Let $m : E \rightarrow \mathbb{R}_+$ be a modular function and define $f(A) = g(m(A))$ where g is concave. Then f is submodular.

Example: Lovász extension of concave over modular

- Let $m : E \rightarrow \mathbb{R}_+$ be a modular function and define $f(A) = g(m(A))$ where g is concave. Then f is submodular.
- Let $M_j = \sum_{i=1}^j m(e_i)$

Example: Lovász extension of concave over modular

- Let $m : E \rightarrow \mathbb{R}_+$ be a modular function and define $f(A) = g(m(A))$ where g is concave. Then f is submodular.
- Let $M_j = \sum_{i=1}^j m(e_i)$
- $\tilde{f}(w)$ is given as

$$\tilde{f}(w) = \sum_{i=1}^m w(e_i) (g(M_i) - g(M_{i-1})) \quad (15.110)$$

Example: Lovász extension of concave over modular

- Let $m : E \rightarrow \mathbb{R}_+$ be a modular function and define $f(A) = g(m(A))$ where g is concave. Then f is submodular.
- Let $M_j = \sum_{i=1}^j m(e_i)$
- $\tilde{f}(w)$ is given as

$$\tilde{f}(w) = \sum_{i=1}^m w(e_i) (g(M_i) - g(M_{i-1})) \quad (15.110)$$

- And if $m(A) = |A|$, we get

$$\tilde{f}(w) = \sum_{i=1}^m w(e_i) (g(i) - g(i-1)) \quad (15.111)$$

Example: Lovász extension and cut functions

- **Cut Function:** Given a non-negative weighted graph $G = (V, E, m)$ where $m : E \rightarrow \mathbb{R}_+$ is a modular function over the edges, we know from Lecture 2 that $f : 2^V \rightarrow \mathbb{R}_+$ with $f(X) = m(\Gamma(X))$ where $\Gamma(X) = \{(u, v) \mid (u, v) \in E, u \in X, v \in V \setminus X\}$ is non-monotone submodular.

Example: Lovász extension and cut functions

- **Cut Function:** Given a non-negative weighted graph $G = (V, E, m)$ where $m : E \rightarrow \mathbb{R}_+$ is a modular function over the edges, we know from Lecture 2 that $f : 2^V \rightarrow \mathbb{R}_+$ with $f(X) = m(\Gamma(X))$ where $\Gamma(X) = \{(u, v) | (u, v) \in E, u \in X, v \in V \setminus X\}$ is non-monotone submodular.
- Simple way to write it, with $m_{ij} = m((i, j))$:

$$f(X) = \sum_{i \in X, j \in V \setminus X} m_{ij} \quad (15.112)$$

Example: Lovász extension and cut functions

- **Cut Function:** Given a non-negative weighted graph $G = (V, E, m)$ where $m : E \rightarrow \mathbb{R}_+$ is a modular function over the edges, we know from Lecture 2 that $f : 2^V \rightarrow \mathbb{R}_+$ with $f(X) = m(\Gamma(X))$ where $\Gamma(X) = \{(u, v) | (u, v) \in E, u \in X, v \in V \setminus X\}$ is non-monotone submodular.
- Simple way to write it, with $m_{ij} = m((i, j))$:

$$f(X) = \sum_{i \in X, j \in V \setminus X} m_{ij} \quad (15.112)$$

- **Exercise:** show that Lovász extension of graph cut may be written as:

$$\tilde{f}(w) = \sum_{i, j \in V} m_{ij} \max \{(w_i - w_j), 0\} \quad (15.113)$$

where elements are ordered as usual, $w_1 \geq w_2 \geq \dots \geq w_n$.

Example: Lovász extension and cut functions

- **Cut Function:** Given a non-negative weighted graph $G = (V, E, m)$ where $m : E \rightarrow \mathbb{R}_+$ is a modular function over the edges, we know from Lecture 2 that $f : 2^V \rightarrow \mathbb{R}_+$ with $f(X) = m(\Gamma(X))$ where $\Gamma(X) = \{(u, v) | (u, v) \in E, u \in X, v \in V \setminus X\}$ is non-monotone submodular.
- Simple way to write it, with $m_{ij} = m((i, j))$:

$$f(X) = \sum_{i \in X, j \in V \setminus X} m_{ij} \quad (15.112)$$

- **Exercise:** show that Lovász extension of graph cut may be written as:

$$\tilde{f}(w) = \sum_{i, j \in V} m_{ij} \max \{(w_i - w_j), 0\} \quad (15.113)$$

where elements are ordered as usual, $w_1 \geq w_2 \geq \dots \geq w_n$.

- This is also a form of “total variation”

A few more Lovász extension examples

Some additional submodular functions and their Lovász extensions, where $w(e_1) \geq w(e_2) \geq \dots \geq w(e_m) \geq 0$. Let $W_k \triangleq \sum_{i=1}^k w(e_i)$.

$f(A)$	$\tilde{f}(w)$
$ A $	$\ w\ _1$
$\min(A , 1)$	$\ w\ _\infty$
$\min(A , 1) - \max(A - m + 1, 0)$	$\ w\ _\infty - \min_i w_i$
$\min(A , k)$	W_k
$\min(A , k) - \max(A - (n - k) + 1, 1)$	$2W_k - W_m$
$\min(A , E \setminus A)$	$2W_{\lfloor m/2 \rfloor} - W_m$

(thanks to K. Narayanan).

Supervised And Unsupervised Machine Learning

- Given training data $\mathcal{D} = \{(x_i, y_i)\}_{i=1}^m$ with $(x_i, y_i) \in \mathbb{R}^n \times \mathbb{R}$, perform the following risk minimization problem:

$$\min_{w \in \mathbb{R}^n} \frac{1}{m} \sum_{i=1}^m \ell(y_i, w^\top x_i) + \lambda \Omega(w), \quad (15.114)$$

where $\ell(\cdot)$ is a loss function (e.g., squared error) and $\Omega(w)$ is a norm.

- When data has multiple responses $(x_i, y_i) \in \mathbb{R}^n \times \mathbb{R}^k$, learning becomes:

$$\min_{w^1, \dots, w^k \in \mathbb{R}^n} \sum_{j=1}^k \frac{1}{m} \sum_{i=1}^m \ell(y_i^k, (w^k)^\top x_i) + \lambda \Omega(w^k), \quad (15.115)$$

- When data has multiple responses only that are observed, $(y_i) \in \mathbb{R}^k$ we get dictionary learning (Krause & Guestrin, Das & Kempe):

$$\min_{x_1, \dots, x_m} \min_{w^1, \dots, w^k \in \mathbb{R}^n} \sum_{j=1}^k \frac{1}{m} \sum_{i=1}^m \ell(y_i^k, (w^k)^\top x_i) + \lambda \Omega(w^k), \quad (15.116)$$

Norms, sparse norms, and computer vision

- Common norms include p -norm $\Omega(w) = \|w\|_p = (\sum_{i=1}^p w_i^p)^{1/p}$

Norms, sparse norms, and computer vision

- Common norms include p -norm $\Omega(w) = \|w\|_p = (\sum_{i=1}^p w_i^p)^{1/p}$
- 1-norm promotes sparsity (prefer solutions with zero entries).

Norms, sparse norms, and computer vision

- Common norms include p -norm $\Omega(w) = \|w\|_p = (\sum_{i=1}^p w_i^p)^{1/p}$
- 1-norm promotes sparsity (prefer solutions with zero entries).
- Image denoising, **total variation** is useful, norm takes form:

$$\Omega(w) = \sum_{i=2}^N |w_i - w_{i-1}| \quad (15.117)$$

Norms, sparse norms, and computer vision

- Common norms include p -norm $\Omega(w) = \|w\|_p = (\sum_{i=1}^p w_i^p)^{1/p}$
- 1-norm promotes sparsity (prefer solutions with zero entries).
- Image denoising, **total variation** is useful, norm takes form:

$$\Omega(w) = \sum_{i=2}^N |w_i - w_{i-1}| \quad (15.117)$$

- Points of difference should be “sparse” (frequently zero).



(Rodriguez,
2009)

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.
- With $\|w\|_0$ or its relaxation, each non-zero element has equal degree of penalty. Penalties do not interact.

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.
- With $\|w\|_0$ or its relaxation, each non-zero element has equal degree of penalty. Penalties do not interact.
- Given submodular function $f : 2^V \rightarrow \mathbb{R}_+$, $f(\text{supp}(w))$ measures the “complexity” of the non-zero pattern of w ; can have more non-zero values if they cooperate (via f) with other non-zero values.

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.
- With $\|w\|_0$ or its relaxation, each non-zero element has equal degree of penalty. Penalties do not interact.
- Given submodular function $f : 2^V \rightarrow \mathbb{R}_+$, $f(\text{supp}(w))$ measures the “complexity” of the non-zero pattern of w ; can have more non-zero values if they cooperate (via f) with other non-zero values.
- $f(\text{supp}(w))$ is hard to optimize, but its convex envelope $\tilde{f}(\|w\|)$ (i.e., largest convex under-estimator of $f(\text{supp}(w))$) is obtained via the Lovász-extension \tilde{f} of f (Vondrák 2007, Bach 2010).

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.
- With $\|w\|_0$ or its relaxation, each non-zero element has equal degree of penalty. Penalties do not interact.
- Given submodular function $f : 2^V \rightarrow \mathbb{R}_+$, $f(\text{supp}(w))$ measures the “complexity” of the non-zero pattern of w ; can have more non-zero values if they cooperate (via f) with other non-zero values.
- $f(\text{supp}(w))$ is hard to optimize, but its convex envelope $\tilde{f}(\|w\|)$ (i.e., largest convex under-estimator of $f(\text{supp}(w))$) is obtained via the Lovász-extension \tilde{f} of f (Vondrák 2007, Bach 2010).
- Submodular functions thus parameterize structured convex sparse norms via the Lovász-extension!

Submodular parameterization of a sparse convex norm

- Prefer convex norms since they can be solved.
- For $w \in \mathbb{R}^V$, $\text{supp}(w) \in \{0, 1\}^V$ has $\text{supp}(w)(v) = 1$ iff $w(v) > 0$
- Desirable sparse norm: count the non-zeros, $\|w\|_0 = \mathbf{1}^\top \text{supp}(w)$.
- Using $\Omega(w) = \|w\|_0$ is NP-hard, instead we often optimize tightest convex relaxation, $\|w\|_1$ which is the convex envelope.
- With $\|w\|_0$ or its relaxation, each non-zero element has equal degree of penalty. Penalties do not interact.
- Given submodular function $f : 2^V \rightarrow \mathbb{R}_+$, $f(\text{supp}(w))$ measures the “complexity” of the non-zero pattern of w ; can have more non-zero values if they cooperate (via f) with other non-zero values.
- $f(\text{supp}(w))$ is hard to optimize, but it's convex envelope $\tilde{f}(\|w\|)$ (i.e., largest convex under-estimator of $f(\text{supp}(w))$) is obtained via the Lovász-extension \tilde{f} of f (Vondrák 2007, Bach 2010).
- Submodular functions thus parameterize structured convex sparse norms via the Lovász-extension!
- Ex: total variation is Lovász-ext. of graph cut, but \exists many more!

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).
- Simple example. The Lovász extension of the modular function $f(A) = |A|$ is the ℓ_1 norm, and the Lovász extension of the modular function $f(A) = m(A)$ is the weighted ℓ_1 norm.

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).
- Simple example. The Lovász extension of the modular function $f(A) = |A|$ is the ℓ_1 norm, and the Lovász extension of the modular function $f(A) = m(A)$ is the weighted ℓ_1 norm.
- With more general submodular functions, one can generate a large and interesting variety of norms, all of which have polyhedral contours (unlike, say, something like the ℓ_2 norm).

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).
- Simple example. The Lovász extension of the modular function $f(A) = |A|$ is the ℓ_1 norm, and the Lovász extension of the modular function $f(A) = m(A)$ is the weighted ℓ_1 norm.
- With more general submodular functions, one can generate a large and interesting variety of norms, all of which have polyhedral contours (unlike, say, something like the ℓ_2 norm).
- Hence, not all norms come from the Lovász extension of some submodular function.

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).
- Simple example. The Lovász extension of the modular function $f(A) = |A|$ is the ℓ_1 norm, and the Lovász extension of the modular function $f(A) = m(A)$ is the weighted ℓ_1 norm.
- With more general submodular functions, one can generate a large and interesting variety of norms, all of which have polyhedral contours (unlike, say, something like the ℓ_2 norm).
- Hence, not all norms come from the Lovász extension of some submodular function.
- Similarly, not all convex functions are the Lovász extension of some submodular function.

Lovász extension and norms

- Using Lovász extension to define various norms of the form $\|w\|_{\tilde{f}} = \tilde{f}(|w|)$, renders the function symmetric about all orthants (i.e., $\|w\|_{\tilde{f}} = \|b \odot w\|_{\tilde{f}}$ where $b \in \{-1, 1\}^m$ and \odot is element-wise multiplication).
- Simple example. The Lovász extension of the modular function $f(A) = |A|$ is the ℓ_1 norm, and the Lovász extension of the modular function $f(A) = m(A)$ is the weighted ℓ_1 norm.
- With more general submodular functions, one can generate a large and interesting variety of norms, all of which have polyhedral contours (unlike, say, something like the ℓ_2 norm).
- Hence, not all norms come from the Lovász extension of some submodular function.
- Similarly, not all convex functions are the Lovász extension of some submodular function.
- [Bach-2011](#) has a complete discussion of this.