

# Submodular Functions, Optimization, and Applications to Machine Learning

— Spring Quarter, Lecture 8 —

[http://www.ee.washington.edu/people/faculty/bilmes/classes/ee563\\_spring\\_2018/](http://www.ee.washington.edu/people/faculty/bilmes/classes/ee563_spring_2018/)

Prof. Jeff Bilmes

University of Washington, Seattle  
Department of Electrical Engineering  
<http://melodi.ee.washington.edu/~bilmes>

April 18th, 2018



$$f(A) + f(B) \geq f(A \cup B) + f(A \cap B)$$

$$-f(A) + 2f(C) + f(B) \quad -f(A) + f(C) + f(B) \quad -f(A \cap B)$$



# Cumulative Outstanding Reading

- Read chapter 1 from Fujishige's book.
- Read chapter 2 from Fujishige's book.

# Announcements, Assignments, and Reminders

- If you have any questions about anything, please ask them via our discussion board ([https://canvas.uw.edu/courses/1216339/discussion\\_topics](https://canvas.uw.edu/courses/1216339/discussion_topics)).

# Class Road Map - EE563

- L1(3/26): Motivation, Applications, & Basic Definitions,
- L2(3/28): Machine Learning Apps (diversity, complexity, parameter, learning target, surrogate).
- L3(4/2): Info theory exs, more apps, definitions, graph/combinatorial examples
- L4(4/4): Graph and Combinatorial Examples, Matrix Rank, Examples and Properties, visualizations
- L5(4/9): More Examples/Properties/ Other Submodular Defs., Independence,
- L6(4/11): Matroids, Matroid Examples, Matroid Rank, Partition/Laminar Matroids
- L7(4/16): Laminar Matroids, System of Distinct Reprs, Transversals, Transversal Matroid, Matroid Representation, Dual Matroids
- L8(4/18): Dual Matroids, Other Matroid Properties, Combinatorial Geometries, Matroids and Greedy.
- L9(4/23): Polyhedra, Matroid Polytopes, Matroids  $\rightarrow$  Polymatroids
- L10(4/29): Matroids  $\rightarrow$  Polymatroids, Polymatroids, Polymatroids and Greedy,
- L11(4/30):
- L12(5/2):
- L13(5/7):
- L14(5/9):
- L15(5/14):
- L16(5/16):
- L17(5/21):
- L18(5/23):
- L-(5/28): Memorial Day (holiday)
- L19(5/30):
- L21(6/4): Final Presentations maximization.

Last day of instruction, June 1st. Finals Week: June 2-8, 2018.

# System of Distinct Representatives

- Let  $(V, \mathcal{V})$  be a set system (i.e.,  $\mathcal{V} = (V_i : i \in I)$  where  $V_i \subseteq V$  for all  $i$ ), and  $I$  is an index set. Hence,  $|I| = |\mathcal{V}|$ .
- A family  $(v_i : i \in I)$  with  $v_i \in V$  is said to be a **system of distinct representatives** of  $\mathcal{V}$  if  $\exists$  a bijection  $\pi : I \leftrightarrow I$  such that  $v_i \in V_{\pi(i)}$  and  $v_i \neq v_j$  for all  $i \neq j$ .
- In a system of distinct representatives, there **is** a requirement for the representatives to be distinct. We can re-state (and rename) this as a:

## Definition 8.2.1 (transversal)

Given a set system  $(V, \mathcal{V})$  and index set  $I$  for  $\mathcal{V}$  as defined above, a set  $T \subseteq V$  is a **transversal** of  $\mathcal{V}$  if there is a bijection  $\pi : T \leftrightarrow I$  such that

$$x \in V_{\pi(x)} \text{ for all } x \in T \quad (8.2)$$

- Note that due to  $\pi : T \leftrightarrow I$  being a bijection, all of  $I$  and  $T$  are “covered” (so this makes things distinct automatically).

# When do transversals exist?

- As we saw, a transversal might not always exist. How to tell?
- Given a set system  $(V, \mathcal{V})$  with  $\mathcal{V} = (V_i : i \in I)$ , and  $V_i \subseteq V$  for all  $i$ . Then, for any  $J \subseteq I$ , let

$$V(J) = \cup_{j \in J} V_j \quad (8.2)$$

so  $|V(J)| : 2^I \rightarrow \mathbb{Z}_+$  is the set cover func. (we know is submodular).

- We have

## Theorem 8.2.1 (Hall's theorem)

*Given a set system  $(V, \mathcal{V})$ , the family of subsets  $\mathcal{V} = (V_i : i \in I)$  has a transversal  $(v_i : i \in I)$  iff for all  $J \subseteq I$*

$$|V(J)| \geq |J| \quad (8.3)$$


# When do transversals exist?

- As we saw, a transversal might not always exist. How to tell?
- Given a set system  $(V, \mathcal{V})$  with  $\mathcal{V} = (V_i : i \in I)$ , and  $V_i \subseteq V$  for all  $i$ . Then, for any  $J \subseteq I$ , let

$$V(J) = \cup_{j \in J} V_j \quad (8.2)$$

so  $|V(J)| : 2^I \rightarrow \mathbb{Z}_+$  is the set cover func. (we know is submodular).

- Hall's theorem ( $\forall J \subseteq I, |V(J)| \geq |J|$ ) as a bipartite graph.




# When do transversals exist?

- As we saw, a transversal might not always exist. How to tell?
- Given a set system  $(V, \mathcal{V})$  with  $\mathcal{V} = (V_i : i \in I)$ , and  $V_i \subseteq V$  for all  $i$ . Then, for any  $J \subseteq I$ , let

$$V(J) = \cup_{j \in J} V_j \quad (8.2)$$

so  $|V(J)| : 2^I \rightarrow \mathbb{Z}_+$  is the set cover func. (we know is submodular).

- Hall's theorem ( $\forall J \subseteq I, |V(J)| \geq |J|$ ) as a bipartite graph.





## When do transversals exist?

- As we saw, a transversal might not always exist. How to tell?
- Given a set system  $(V, \mathcal{V})$  with  $\mathcal{V} = (V_i : i \in I)$ , and  $V_i \subseteq V$  for all  $i$ . Then, for any  $J \subseteq I$ , let

$$V(J) = \cup_{j \in J} V_j \quad (8.2)$$

so  $|V(J)| : 2^I \rightarrow \mathbb{Z}_+$  is the set cover func. (we know is submodular).

- Moreover, we have

### Theorem 8.2.2 (Rado's theorem (1942))

*If  $M = (V, r)$  is a matroid on  $V$  with rank function  $r$ , then the family of subsets  $(V_i : i \in I)$  of  $V$  has a transversal  $(v_i : i \in I)$  that is independent in  $M$  iff for all  $J \subseteq I$*

$$r(V(J)) \geq |J| \quad (8.4)$$

- Note, a transversal  $T$  independent in  $M$  means that  $r(T) = |T|$ .

# Application's of Hall's theorem

- Consider a set of jobs  $I$  and a set of applicants  $V$  to the jobs. If an applicant  $v \in V$  is qualified for job  $i \in I$ , we add edge  $(v, i)$  to the bipartite graph  $G = (V, I, E)$ .

# Application's of Hall's theorem

- Consider a set of jobs  $I$  and a set of applicants  $V$  to the jobs. If an applicant  $v \in V$  is qualified for job  $i \in I$ , we add edge  $(v, i)$  to the bipartite graph  $G = (V, I, E)$ .
- We wish all jobs to be filled, and hence Hall's condition  $(\forall J \subseteq I, |V(J)| \geq |J|)$  is a necessary and sufficient condition for this to be possible.

# Application's of Hall's theorem

- Consider a set of jobs  $I$  and a set of applicants  $V$  to the jobs. If an applicant  $v \in V$  is qualified for job  $i \in I$ , we add edge  $(v, i)$  to the bipartite graph  $G = (V, I, E)$ .
- We wish all jobs to be filled, and hence Hall's condition  $(\forall J \subseteq I, |V(J)| \geq |J|)$  is a necessary and sufficient condition for this to be possible.
- Note if  $|V| = |I|$ , then Hall's theorem is the Marriage Theorem (Frobenius 1917), where an edge  $(v, i)$  in the graph indicate compatibility between two individuals  $v \in V$  and  $i \in I$  coming from two separate groups  $V$  and  $I$ .

# Application's of Hall's theorem

- Consider a set of jobs  $I$  and a set of applicants  $V$  to the jobs. If an applicant  $v \in V$  is qualified for job  $i \in I$ , we add edge  $(v, i)$  to the bipartite graph  $G = (V, I, E)$ .
- We wish all jobs to be filled, and hence Hall's condition  $(\forall J \subseteq I, |V(J)| \geq |J|)$  is a necessary and sufficient condition for this to be possible.
- Note if  $|V| = |I|$ , then Hall's theorem is the Marriage Theorem (Frobenius 1917), where an edge  $(v, i)$  in the graph indicate compatibility between two individuals  $v \in V$  and  $i \in I$  coming from two separate groups  $V$  and  $I$ .
- If  $\forall J \subseteq I, |V(J)| \geq |J|$ , then all individuals in each group can be matched with a compatible mate.

# More general conditions for existence of transversals

## Theorem 8.2.1 (Polymatroid transversal theorem)

If  $\mathcal{V} = (V_i : i \in I)$  is a finite family of non-empty subsets of  $V$ , and  $f : 2^V \rightarrow \mathbb{Z}_+$  is a non-negative, integral, monotone non-decreasing, and submodular function, then  $\mathcal{V}$  has a system of representatives  $(v_i : i \in I)$  such that

$$f(\cup_{i \in J} \{v_i\}) \geq |J| \text{ for all } J \subseteq I \quad (8.2)$$

if and only if

$$f(V(J)) \geq |J| \text{ for all } J \subseteq I \quad (8.3)$$

- Given Theorem ??, we immediately get Theorem 8.2.1 by taking  $f(S) = |S|$  for  $S \subseteq V$ .
- We get Theorem ?? by taking  $f(S) = r(S)$  for  $S \subseteq V$ , the rank function of the matroid.

# Review from Lecture 6

The next frame comes from lecture 6.

# Matroids, other definitions using matroid rank $r : 2^V \rightarrow \mathbb{Z}_+$

## Definition 8.3.3 (closed/flat/subspace)

A subset  $A \subseteq E$  is **closed** (equivalently, a **flat** or a **subspace**) of matroid  $M$  if for all  $x \in E \setminus A$ ,  $r(A \cup \{x\}) = r(A) + 1$ .

Definition: A **hyperplane** is a flat of rank  $r(M) - 1$ .

## Definition 8.3.4 (closure)

Given  $A \subseteq E$ , the **closure** (or **span**) of  $A$ , is defined by  $\text{span}(A) = \{b \in E : r(A \cup \{b\}) = r(A)\}$ .

Therefore, a closed set  $A$  has  $\text{span}(A) = A$ .

## Definition 8.3.5 (circuit)

A subset  $A \subseteq E$  is **circuit** or a **cycle** if it is an inclusionwise-minimal dependent set (i.e., if  $r(A) < |A|$  and for any  $a \in A$ ,  $r(A \setminus \{a\}) = |A| - 1$ ).



# Spanning Sets

- We have the following definitions:

# Spanning Sets

- We have the following definitions:

## Definition 8.3.1 (spanning set of a set)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , and a set  $Y \subseteq V$ , then any set  $X \subseteq Y$  such that  $r(X) = r(Y)$  is called a **spanning set** of  $Y$ .

# Spanning Sets

- We have the following definitions:

## Definition 8.3.1 (spanning set of a set)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , and a set  $Y \subseteq V$ , then any set  $X \subseteq Y$  such that  $r(X) = r(Y)$  is called a **spanning set** of  $Y$ .

## Definition 8.3.2 (spanning set of a matroid)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , any set  $A \subseteq V$  such that  $r(A) = r(V)$  is called a **spanning set** of the matroid.

# Spanning Sets

- We have the following definitions:

## Definition 8.3.1 (spanning set of a set)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , and a set  $Y \subseteq V$ , then any set  $X \subseteq Y$  such that  $r(X) = r(Y)$  is called a **spanning set** of  $Y$ .

## Definition 8.3.2 (spanning set of a matroid)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , any set  $A \subseteq V$  such that  $r(A) = r(V)$  is called a **spanning set** of the matroid.

- A base of a matroid is a minimal spanning set (and it is independent) but supersets of a base are also spanning.

# Spanning Sets

- We have the following definitions:

## Definition 8.3.1 (spanning set of a set)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , and a set  $Y \subseteq V$ , then any set  $X \subseteq Y$  such that  $r(X) = r(Y)$  is called a **spanning set** of  $Y$ .

## Definition 8.3.2 (spanning set of a matroid)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , any set  $A \subseteq V$  such that  $r(A) = r(V)$  is called a **spanning set** of the matroid.

- A base of a matroid is a minimal spanning set (and it is independent) but supersets of a base are also spanning.
- $V$  is always trivially spanning.

# Spanning Sets

- We have the following definitions:

## Definition 8.3.1 (spanning set of a set)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , and a set  $Y \subseteq V$ , then any set  $X \subseteq Y$  such that  $r(X) = r(Y)$  is called a **spanning set** of  $Y$ .

## Definition 8.3.2 (spanning set of a matroid)

Given a matroid  $\mathcal{M} = (V, \mathcal{I})$ , any set  $A \subseteq V$  such that  $r(A) = r(V)$  is called a **spanning set** of the matroid.

- A base of a matroid is a minimal spanning set (and it is independent) but supersets of a base are also spanning.
- $V$  is always trivially spanning.
- Consider the terminology: “spanning tree in a graph”, comes from spanning in a matroid sense.

# Dual of a Matroid

- Given a matroid  $M = (V, \mathcal{I})$ , a dual matroid  $M^* = (V, \mathcal{I}^*)$  can be defined on the same ground set  $V$ , but using a **very different** set of independent sets  $\mathcal{I}^*$ .

# Dual of a Matroid

- Given a matroid  $M = (V, \mathcal{I})$ , a dual matroid  $M^* = (V, \mathcal{I}^*)$  can be defined on the same ground set  $V$ , but using a **very different** set of independent sets  $\mathcal{I}^*$ .
- We define the set of sets  $\mathcal{I}^*$  for  $M^*$  as follows:

$$\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\} \quad (8.1)$$

$$= \{V \setminus S : S \subseteq V \text{ is a spanning set of } M\} \quad (8.2)$$

i.e.,  $\mathcal{I}^*$  are complements of spanning sets of  $M$ .



# Dual of a Matroid

- Given a matroid  $M = (V, \mathcal{I})$ , a dual matroid  $M^* = (V, \mathcal{I}^*)$  can be defined on the same ground set  $V$ , but using a **very different** set of independent sets  $\mathcal{I}^*$ .
- We define the set of sets  $\mathcal{I}^*$  for  $M^*$  as follows:

$$\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\} \quad (8.1)$$

$$= \{V \setminus S : S \subseteq V \text{ is a spanning set of } M\} \quad (8.2)$$

i.e.,  $\mathcal{I}^*$  are complements of spanning sets of  $M$ .

- That is, a set  $A$  is independent in the dual matroid  $M^*$  if removal of  $A$  from  $V$  does not decrease the rank in  $M$ :

$$\mathcal{I}^* = \{A \subseteq V : \text{rank}_M(V \setminus A) = \text{rank}_M(V)\} \quad (8.3)$$

# Dual of a Matroid

- Given a matroid  $M = (V, \mathcal{I})$ , a dual matroid  $M^* = (V, \mathcal{I}^*)$  can be defined on the same ground set  $V$ , but using a **very different** set of independent sets  $\mathcal{I}^*$ .
- We define the set of sets  $\mathcal{I}^*$  for  $M^*$  as follows:

$$\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\} \quad (8.1)$$

$$= \{V \setminus S : S \subseteq V \text{ is a spanning set of } M\} \quad (8.2)$$

i.e.,  $\mathcal{I}^*$  are complements of spanning sets of  $M$ .

- That is, a set  $A$  is independent in the dual matroid  $M^*$  if removal of  $A$  from  $V$  does not decrease the rank in  $M$ :

$$\mathcal{I}^* = \{A \subseteq V : \text{rank}_M(V \setminus A) = \text{rank}_M(V)\} \quad (8.3)$$

- In other words, a set  $A \subseteq V$  is independent in the dual  $M^*$  (i.e.,  $A \in \mathcal{I}^*$ ) if  $A$ 's complement is spanning in  $M$  (residual  $V \setminus A$  must contain a base in  $M$ ).

# Dual of a Matroid

- Given a matroid  $M = (V, \mathcal{I})$ , a dual matroid  $M^* = (V, \mathcal{I}^*)$  can be defined on the same ground set  $V$ , but using a **very different** set of independent sets  $\mathcal{I}^*$ .
- We define the set of sets  $\mathcal{I}^*$  for  $M^*$  as follows:

$$\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\} \quad (8.1)$$

$$= \{V \setminus S : S \subseteq V \text{ is a spanning set of } M\} \quad (8.2)$$

i.e.,  $\mathcal{I}^*$  are complements of spanning sets of  $M$ .

- That is, a set  $A$  is independent in the dual matroid  $M^*$  if removal of  $A$  from  $V$  does not decrease the rank in  $M$ :

$$\mathcal{I}^* = \{A \subseteq V : \text{rank}_M(V \setminus A) = \text{rank}_M(V)\} \quad (8.3)$$

- In other words, a set  $A \subseteq V$  is independent in the dual  $M^*$  (i.e.,  $A \in \mathcal{I}^*$ ) if  $A$ 's complement is spanning in  $M$  (residual  $V \setminus A$  must contain a base in  $M$ ).
- Dual of the dual:** Note, we have that  $(M^*)^* = M$ .

# Dual of a Matroid: Bases

- The smallest spanning sets are bases.

# Dual of a Matroid: Bases

- The smallest spanning sets are bases. Hence, a base  $B$  of  $M$  (where  $B = V \setminus B^*$  is as small as possible while still spanning) is the complement of a base  $B^*$  of  $M^*$  (where  $B^* = V \setminus B$  is as large as possible while still being independent).

# Dual of a Matroid: Bases

- The smallest spanning sets are bases. Hence, a base  $B$  of  $M$  (where  $B = V \setminus B^*$  is as small as possible while still spanning) is the complement of a base  $B^*$  of  $M^*$  (where  $B^* = V \setminus B$  is as large as possible while still being independent).
- In fact, we have that

# Dual of a Matroid: Bases

- The smallest spanning sets are bases. Hence, a base  $B$  of  $M$  (where  $B = V \setminus B^*$  is as small as possible while still spanning) is the complement of a base  $B^*$  of  $M^*$  (where  $B^* = V \setminus B$  is as large as possible while still being independent).
- In fact, we have that

## Theorem 8.3.3 (Dual matroid bases)

Let  $M = (V, \mathcal{I})$  be a matroid and  $\mathcal{B}(M)$  be the set of bases of  $M$ . Then define

$$\mathcal{B}^*(M) = \{V \setminus B : B \in \mathcal{B}(M)\}. \quad (8.4)$$

Then  $\mathcal{B}^*(M)$  is the set of basis of  $M^*$  (that is,  $\mathcal{B}^*(M) = \mathcal{B}(M^*)$ ).

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .



# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

## Proposition 8.3.4 (from Oxley 2011)

*Let  $M = (V, \mathcal{I})$  be a matroid, and let  $X \subseteq V$ . Then*

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

## Proposition 8.3.4 (from Oxley 2011)

Let  $M = (V, \mathcal{I})$  be a matroid, and let  $X \subseteq V$ . Then

- 1  $X$  is independent in  $M$  iff  $V \setminus X$  is cospanning in  $M$  (spanning in  $M^*$ ).

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

## Proposition 8.3.4 (from Oxley 2011)

Let  $M = (V, \mathcal{I})$  be a matroid, and let  $X \subseteq V$ . Then

- 1  $X$  is independent in  $M$  iff  $V \setminus X$  is cospanning in  $M$  (spanning in  $M^*$ ).
- 2  $X$  is spanning in  $M$  iff  $V \setminus X$  is coindependent in  $M$  (independent in  $M^*$ ).

# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

## Proposition 8.3.4 (from Oxley 2011)

Let  $M = (V, \mathcal{I})$  be a matroid, and let  $X \subseteq V$ . Then

- 1  $X$  is independent in  $M$  iff  $V \setminus X$  is cospanning in  $M$  (spanning in  $M^*$ ).
- 2  $X$  is spanning in  $M$  iff  $V \setminus X$  is coindependent in  $M$  (independent in  $M^*$ ).
- 3  $X$  is a hyperplane in  $M$  iff  $V \setminus X$  is a cocircuit in  $M$  (circuit in  $M^*$ ).



# An exercise in duality Terminology

- $\mathcal{B}^*(M)$ , the bases of  $M^*$ , are called **cobases** of  $M$ .
- The circuits of  $M^*$  are called **cocircuits** of  $M$ .
- The hyperplanes of  $M^*$  are called **cohyperplanes** of  $M$ .
- The independent sets of  $M^*$  are called **coindependent** sets of  $M$ .
- The spanning sets of  $M^*$  are called **cospanning** sets of  $M$ .

## Proposition 8.3.4 (from Oxley 2011)

Let  $M = (V, \mathcal{I})$  be a matroid, and let  $X \subseteq V$ . Then

- 1  $X$  is independent in  $M$  iff  $V \setminus X$  is cospanning in  $M$  (spanning in  $M^*$ ).
- 2  $X$  is spanning in  $M$  iff  $V \setminus X$  is coindependent in  $M$  (independent in  $M^*$ ).
- 3  $X$  is a hyperplane in  $M$  iff  $V \setminus X$  is a cocircuit in  $M$  (circuit in  $M^*$ ).
- 4  $X$  is a circuit in  $M$  iff  $V \setminus X$  is a cohyperplane in  $M$  (hyperplane in  $M^*$ ).

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).
- A **cut** in a graph  $G$  is a set of edges, the removal of which increases the number of connected components. I.e.,  $X \subseteq E(G)$  is a cut in  $G$  if  $k(G) < k(G \setminus X)$ .

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).
- A **cut** in a graph  $G$  is a set of edges, the removal of which increases the number of connected components. I.e.,  $X \subseteq E(G)$  is a cut in  $G$  if  $k(G) < k(G \setminus X)$ .
- A **minimal cut** in  $G$  is a cut  $X \subseteq E(G)$  such that  $X \setminus \{x\}$  is not a cut for any  $x \in X$ .

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).
- A **cut** in a graph  $G$  is a set of edges, the removal of which increases the number of connected components. I.e.,  $X \subseteq E(G)$  is a cut in  $G$  if  $k(G) < k(G \setminus X)$ .
- A **minimal cut** in  $G$  is a cut  $X \subseteq E(G)$  such that  $X \setminus \{x\}$  is not a cut for any  $x \in X$ .
- A **cocycle** (cocircuit) in a graphic matroid is a minimal graph cut.

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).
- A **cut** in a graph  $G$  is a set of edges, the removal of which increases the number of connected components. I.e.,  $X \subseteq E(G)$  is a cut in  $G$  if  $k(G) < k(G \setminus X)$ .
- A **minimal cut** in  $G$  is a cut  $X \subseteq E(G)$  such that  $X \setminus \{x\}$  is not a cut for any  $x \in X$ .
- A **cocycle** (cocircuit) in a graphic matroid is a minimal graph cut.
- A mincut is a circuit in the dual “cocycle” (or “cut”) matroid.

## Example duality: graphic matroid

- Using a graphic/cycle matroid, we can already see how dual matroid concepts demonstrates the extraordinary flexibility and power that a matroid can have.
- Recall, in cycle matroid, a spanning set of  $G$  is any set of edges that are incident to all nodes (i.e., any superset of a spanning forest), a minimal spanning set is a spanning tree (or forest), and a circuit has a nice visual interpretation (a cycle in the graph).
- A **cut** in a graph  $G$  is a set of edges, the removal of which increases the number of connected components. I.e.,  $X \subseteq E(G)$  is a cut in  $G$  if  $k(G) < k(G \setminus X)$ .
- A **minimal cut** in  $G$  is a cut  $X \subseteq E(G)$  such that  $X \setminus \{x\}$  is not a cut for any  $x \in X$ .
- A **cocycle** (cocircuit) in a graphic matroid is a minimal graph cut.
- A mincut is a circuit in the dual “cocycle” (or “cut”) matroid.
- All dependent sets in a cocycle matroid are cuts (i.e., a dependent set is a minimal cut or contains one).



# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$


## Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.


## A graph G




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Minimally spanning in  $M$  (and thus a base (maximally independent) in  $M$ )




Maximally independent in  $M^*$  (thus a base, minimally spanning, in  $M^*$ )




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Minimally spanning in  $M$  (and thus a base (maximally independent) in  $M$ )




Maximally independent in  $M^*$  (thus a base, minimally spanning, in  $M^*$ )




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Independent but not spanning  
in  $M$ , and not closed in  $M$ .




Dependent in  $M^*$  (contains  
a cocycle, is a nonminimal cut)




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Spanning in  $M$ , but not a base, and not independent (has cycles)




Independent in  $M^*$  (does not contain a cut)




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Independent but not spanning  
in  $M$ , and not closed in  $M$ .



Dependent in  $M^*$  (contains  
a cocycle, is a nonminimal cut)







# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

A hyperplane in  $M$ , dependent but not spanning in  $M$




A cycle in  $M^*$  (minimally dependent in  $M^*$ , a cocycle, or a minimal cut)




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

A hyperplane in  $M$ , dependent but not spanning in  $M$




A cycle in  $M^*$  (minimally dependent in  $M^*$ , a cocycle, or a minimal cut)




# Example: cocycle matroid (sometimes “cut matroid”)

- The dual of the cycle matroid is called the cocycle matroid. Recall,  $\mathcal{I}^* = \{A \subseteq V : V \setminus A \text{ is a spanning set of } M\}$
- $\mathcal{I}^*$  consists of all sets of edges the complement of which contains a spanning tree — i.e., an independent set can't consist of edges that, if removed, would render the graph non-spanning.

Cycle Matroid - independent sets have no cycles.



Cocycle matroid, independent sets contain no cuts.



# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

*Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.*

## Proof.

- Since  $V \setminus \emptyset$  is spanning in primal, clearly  $\emptyset \in \mathcal{I}^*$ , so (I1') holds.

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Since  $V \setminus \emptyset$  is spanning in primal, clearly  $\emptyset \in \mathcal{I}^*$ , so (I1') holds.
- Also, if  $I \subseteq J \in \mathcal{I}^*$ , then clearly also  $I \in \mathcal{I}^*$  since if  $V \setminus J$  is spanning in  $M$ , so must  $V \setminus I$ . Therefore, (I2') holds.
- Next, given  $I, J \in \mathcal{I}^*$  with  $|I| < |J|$ , it must be the case that  $\bar{I} = V \setminus I$  and  $\bar{J} = V \setminus J$  are both spanning in  $M$  with  $|\bar{I}| > |\bar{J}|$ .

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Consider  $I, J \in \mathcal{I}^*$  with  $|I| < |J|$ . We need to show that there is some member  $v \in J \setminus I$  such that  $I + v$  is independent in  $M^*$ , which means that  $V \setminus (I + v) = (V \setminus I) \setminus v = \bar{I} - v$  is still spanning in  $M$ . That is, removing  $v$  from  $V \setminus I$  doesn't make  $(V \setminus I) \setminus v$  not spanning in  $M$ .

...

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Consider  $I, J \in \mathcal{I}^*$  with  $|I| < |J|$ . We need to show that there is some member  $v \in J \setminus I$  such that  $I + v$  is independent in  $M^*$ , which means that  $V \setminus (I + v) = (V \setminus I) \setminus v = \bar{I} - v$  is still spanning in  $M$ . That is, removing  $v$  from  $V \setminus I$  doesn't make  $(V \setminus I) \setminus v$  not spanning in  $M$ .
- Since  $V \setminus J$  is spanning in  $M$ ,  $V \setminus J$  contains some base (say  $B_{\bar{J}} \subseteq V \setminus J$ ) of  $M$ . Also,  $V \setminus I$  contains a base of  $M$ , say  $B_{\bar{I}} \subseteq V \setminus I$ .

...

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Consider  $I, J \in \mathcal{I}^*$  with  $|I| < |J|$ . We need to show that there is some member  $v \in J \setminus I$  such that  $I + v$  is independent in  $M^*$ , which means that  $V \setminus (I + v) = (V \setminus I) \setminus v = \bar{I} - v$  is still spanning in  $M$ . That is, removing  $v$  from  $V \setminus I$  doesn't make  $(V \setminus I) \setminus v$  not spanning in  $M$ .
- Since  $V \setminus J$  is spanning in  $M$ ,  $V \setminus J$  contains some base (say  $B_{\bar{J}} \subseteq V \setminus J$ ) of  $M$ . Also,  $V \setminus I$  contains a base of  $M$ , say  $B_{\bar{I}} \subseteq V \setminus I$ .
- Since  $B_{\bar{J}} \setminus I \subseteq V \setminus I$ , and  $B_{\bar{J}} \setminus I$  is independent in  $M$ , we can choose the base  $B_{\bar{I}}$  of  $M$  s.t.  $B_{\bar{J}} \setminus I \subseteq B_{\bar{I}} \subseteq V \setminus I$ .

...



# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Consider  $I, J \in \mathcal{I}^*$  with  $|I| < |J|$ . We need to show that there is some member  $v \in J \setminus I$  such that  $I + v$  is independent in  $M^*$ , which means that  $V \setminus (I + v) = (V \setminus I) \setminus v = \bar{I} - v$  is still spanning in  $M$ . That is, removing  $v$  from  $V \setminus I$  doesn't make  $(V \setminus I) \setminus v$  not spanning in  $M$ .
- Since  $V \setminus J$  is spanning in  $M$ ,  $V \setminus J$  contains some base (say  $B_{\bar{J}} \subseteq V \setminus J$ ) of  $M$ . Also,  $V \setminus I$  contains a base of  $M$ , say  $B_{\bar{I}} \subseteq V \setminus I$ .
- Since  $B_{\bar{J}} \setminus I \subseteq V \setminus I$ , and  $B_{\bar{J}} \setminus I$  is independent in  $M$ , we can choose the base  $B_{\bar{I}}$  of  $M$  s.t.  $B_{\bar{J}} \setminus I \subseteq B_{\bar{I}} \subseteq V \setminus I$ .
- Since  $B_{\bar{J}}$  and  $J$  are disjoint, we have both: 1)  $B_{\bar{J}} \setminus I$  and  $J \setminus I$  are disjoint; and 2)  $B_{\bar{J}} \cap I \subseteq I \setminus J$ . Also note,  $B_{\bar{I}}$  and  $I$  are disjoint. ...

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Now  $J \setminus I \not\subseteq B_{\bar{I}}$ , since otherwise (i.e., assuming  $J \setminus I \subseteq B_{\bar{I}}$ ):

$$|B_{\bar{J}}| = |B_{\bar{J}} \cap I| + |B_{\bar{J}} \setminus I| \quad (8.5)$$

$$\leq |I \setminus J| + |B_{\bar{J}} \setminus I| \quad (8.6)$$

$$< |J \setminus I| + |B_{\bar{J}} \setminus I| \leq |B_{\bar{I}}| \quad (8.7)$$

which is a contradiction. *The last inequality on the right follows since  $J \setminus I \subseteq B_{\bar{I}}$  (by assumption) and  $B_{\bar{J}} \setminus I \subseteq B_{\bar{I}}$  implies that  $(J \setminus I) \cup (B_{\bar{J}} \setminus I) \subseteq B_{\bar{I}}$ , but since  $J$  and  $B_{\bar{J}}$  are disjoint, we have that  $|J \setminus I| + |B_{\bar{J}} \setminus I| \leq |B_{\bar{I}}|$ .*

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Now  $J \setminus I \not\subseteq B_{\bar{I}}$ , since otherwise (i.e., assuming  $J \setminus I \subseteq B_{\bar{I}}$ ):

$$|B_{\bar{J}}| = |B_{\bar{J}} \cap I| + |B_{\bar{J}} \setminus I| \tag{8.5}$$

$$\leq |I \setminus J| + |B_{\bar{J}} \setminus I| \tag{8.6}$$

$$< |J \setminus I| + |B_{\bar{J}} \setminus I| \leq |B_{\bar{I}}| \tag{8.7}$$

which is a contradiction.

- Therefore,  $J \setminus I \not\subseteq B_{\bar{I}}$ , and there is a  $v \in J \setminus I$  s.t.  $v \notin B_{\bar{I}}$ .

...

# The dual of a matroid is (indeed) a matroid

## Theorem 8.3.5

Given matroid  $M = (V, \mathcal{I})$ , let  $M^* = (V, \mathcal{I}^*)$  be as previously defined. Then  $M^*$  is a matroid.

## Proof.

- Now  $J \setminus I \not\subseteq B_{\bar{I}}$ , since otherwise (i.e., assuming  $J \setminus I \subseteq B_{\bar{I}}$ ):

$$|B_{\bar{J}}| = |B_{\bar{J}} \cap I| + |B_{\bar{J}} \setminus I| \quad (8.5)$$

$$\leq |I \setminus J| + |B_{\bar{J}} \setminus I| \quad (8.6)$$

$$< |J \setminus I| + |B_{\bar{J}} \setminus I| \leq |B_{\bar{I}}| \quad (8.7)$$

which is a contradiction.

- Therefore,  $J \setminus I \not\subseteq B_{\bar{I}}$ , and there is a  $v \in J \setminus I$  s.t.  $v \notin B_{\bar{I}}$ .
- So  $B_{\bar{I}}$  is disjoint with  $I \cup \{v\}$ , means  $B_{\bar{I}} \subseteq V \setminus (I \cup \{v\})$ , or  $V \setminus (I \cup \{v\})$  is spanning in  $M$ , and therefore  $I \cup \{v\} \in \mathcal{I}^*$ .



# Matroid Duals and Representability

## Theorem 8.3.6

*Let  $M$  be an  $\mathbb{F}$ -representable matroid (i.e., one that can be represented by a finite sized matrix over field  $\mathbb{F}$ ). Then  $M^*$  is also  $\mathbb{F}$ -representable.*

Hence, for matroids as general as matric matroids, duality does not extend the space of matroids that can be used.

# Matroid Duals and Representability

## Theorem 8.3.6

*Let  $M$  be an  $\mathbb{F}$ -representable matroid (i.e., one that can be represented by a finite sized matrix over field  $\mathbb{F}$ ). Then  $M^*$  is also  $\mathbb{F}$ -representable.*

Hence, for matroids as general as matric matroids, duality does not extend the space of matroids that can be used.

## Theorem 8.3.7

*Let  $M$  be a graphic matroid (i.e., one that can be represented by a graph  $G = (V, E)$ ). Then  $M^*$  is not necessarily also graphic.*

Hence, for graphic matroids, duality can increase the space and power of matroids, and since they are based on a graph, they are relatively easy to use: 1) all cuts are dependent sets; 2) minimal cuts are cycles; 3) bases of a cut are any one edge removed from minimal cuts; 4) independent sets are edges that are not cuts (minimal or otherwise); 5) bases of matroid are maximal non-cuts (non-cut containing edge sets).

# Dual Matroid Rank

## Theorem 8.3.8

The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

- Note, we again immediately see that this is submodular by the properties of submodular functions we saw in lectures 1 and 2. *I.e.,  $|X|$  is modular, complement  $f(V \setminus X)$  is submodular if  $f$  is submodular,  $r_M(V)$  is a constant, and summing submodular functions and a constant preserves submodularity.*

# Dual Matroid Rank

## Theorem 8.3.8

The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

- Note, we again immediately see that this is submodular by the properties of submodular functions we saw in lectures 1 and 2.
- Non-negativity integral follows since  $|X| + r_M(V \setminus X) \geq r_M(X) + r_M(V \setminus X) \geq r_M(V)$ . *The right inequality follows since  $r_M$  is submodular.*



# Dual Matroid Rank

## Theorem 8.3.8

*The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :*

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

- Note, we again immediately see that this is submodular by the properties of submodular functions we saw in lectures 1 and 2.
- Non-negativity integral follows since
$$|X| + r_M(V \setminus X) \geq r_M(X) + r_M(V \setminus X) \geq r_M(V).$$
- Monotone non-decreasing follows since, as  $X$  increases by one,  $|X|$  always increases by 1, while  $r_M(V \setminus X)$  decreases by one or zero.

# Dual Matroid Rank

## Theorem 8.3.8

*The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :*

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

- Note, we again immediately see that this is submodular by the properties of submodular functions we saw in lectures 1 and 2.
- Non-negativity integral follows since  $|X| + r_M(V \setminus X) \geq r_M(X) + r_M(V \setminus X) \geq r_M(V)$ .
- Monotone non-decreasing follows since, as  $X$  increases by one,  $|X|$  always increases by 1, while  $r_M(V \setminus X)$  decreases by one or zero.
- Therefore,  $r_{M^*}$  is the rank function of a matroid. That it is the dual matroid rank function is shown in the next proof.

# Dual Matroid Rank

## Theorem 8.3.8

*The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :*

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

## Proof.

A set  $X$  is independent in  $(V, r_{M^*})$  if and only if

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) = |X| \quad (8.9)$$

...

# Dual Matroid Rank

## Theorem 8.3.8

*The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :*

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

## Proof.

A set  $X$  is independent in  $(V, r_{M^*})$  if and only if

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) = |X| \quad (8.9)$$

or

$$r_M(V \setminus X) = r_M(V) \quad (8.10)$$

...

# Dual Matroid Rank

## Theorem 8.3.8

The rank function  $r_{M^*}$  of the dual matroid  $M^*$  may be specified in terms of the rank  $r_M$  in matroid  $M$  as follows. For  $X \subseteq V$ :

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) \quad (8.8)$$

## Proof.

A set  $X$  is independent in  $(V, r_{M^*})$  if and only if

$$r_{M^*}(X) = |X| + r_M(V \setminus X) - r_M(V) = |X| \quad (8.9)$$

or

$$r_M(V \setminus X) = r_M(V) \quad (8.10)$$

But a subset  $X$  is independent in  $M^*$  only if  $V \setminus X$  is spanning in  $M$  (by the definition of the dual matroid). □

# Matroid restriction/deletion

- Let  $M = (V, \mathcal{I})$  be a matroid and let  $Y \subseteq V$ , then

$$\mathcal{I}_Y = \{Z : Z \subseteq Y, Z \in \mathcal{I}\} \quad (8.11)$$

is such that  $M_Y = (Y, \mathcal{I}_Y)$  is a matroid with rank  $r(M_Y) = r(Y)$ .

# Matroid restriction/deletion

- Let  $M = (V, \mathcal{I})$  be a matroid and let  $Y \subseteq V$ , then

$$\mathcal{I}_Y = \{Z : Z \subseteq Y, Z \in \mathcal{I}\} \quad (8.11)$$

is such that  $M_Y = (Y, \mathcal{I}_Y)$  is a matroid with rank  $r(M_Y) = r(Y)$ .

- This is called the **restriction** of  $M$  to  $Y$ , and is **often written**  $M|Y$ .

# Matroid restriction/deletion

- Let  $M = (V, \mathcal{I})$  be a matroid and let  $Y \subseteq V$ , then

$$\mathcal{I}_Y = \{Z : Z \subseteq Y, Z \in \mathcal{I}\} \quad (8.11)$$

is such that  $M_Y = (Y, \mathcal{I}_Y)$  is a matroid with rank  $r(M_Y) = r(Y)$ .

- This is called the **restriction** of  $M$  to  $Y$ , and is **often written**  $M|Y$ .
- If  $Y = V \setminus X$ , then we have that  $M|Y$  has the form:

$$\mathcal{I}_Y = \{Z : Z \cap X = \emptyset, Z \in \mathcal{I}\} \quad (8.12)$$

is considered a **deletion** of  $X$  from  $M$ , and is **often written**  $M \setminus X$ .



# Matroid restriction/deletion

- Let  $M = (V, \mathcal{I})$  be a matroid and let  $Y \subseteq V$ , then

$$\mathcal{I}_Y = \{Z : Z \subseteq Y, Z \in \mathcal{I}\} \quad (8.11)$$

is such that  $M_Y = (Y, \mathcal{I}_Y)$  is a matroid with rank  $r(M_Y) = r(Y)$ .

- This is called the **restriction** of  $M$  to  $Y$ , and is **often written**  $M|Y$ .
- If  $Y = V \setminus X$ , then we have that  $M|Y$  has the form:

$$\mathcal{I}_Y = \{Z : Z \cap X = \emptyset, Z \in \mathcal{I}\} \quad (8.12)$$

is considered a **deletion** of  $X$  from  $M$ , and is **often written**  $M \setminus X$ .

- Hence,  $M|Y = M \setminus (V \setminus Y)$ , and  $M|(V \setminus X) = M \setminus X$ .

# Matroid restriction/deletion

- Let  $M = (V, \mathcal{I})$  be a matroid and let  $Y \subseteq V$ , then

$$\mathcal{I}_Y = \{Z : Z \subseteq Y, Z \in \mathcal{I}\} \quad (8.11)$$

is such that  $M_Y = (Y, \mathcal{I}_Y)$  is a matroid with rank  $r(M_Y) = r(Y)$ .

- This is called the **restriction** of  $M$  to  $Y$ , and is **often written**  $M|Y$ .
- If  $Y = V \setminus X$ , then we have that  $M|Y$  has the form:

$$\mathcal{I}_Y = \{Z : Z \cap X = \emptyset, Z \in \mathcal{I}\} \quad (8.12)$$

is considered a **deletion** of  $X$  from  $M$ , and is **often written**  $M \setminus X$ .

- Hence,  $M|Y = M \setminus (V \setminus Y)$ , and  $M|(V \setminus X) = M \setminus X$ .
- The rank function is of the same form. I.e.,  $r_Y : 2^Y \rightarrow \mathbb{Z}_+$ , where  $r_Y(Z) = r(Z)$  for  $Z \subseteq Y$ ,  $Y = V \setminus X$ .

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ .  
Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ . Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .
- Let  $Z \subseteq V$  and let  $B_Z$  be a base of  $Z$ . Then a subset  $I \subseteq V \setminus Z$  is independent in  $M/Z$  iff  $I \cup B_Z$  is independent in  $M$ .

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ . Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .
- Let  $Z \subseteq V$  and let  $B_Z$  be a base of  $Z$ . Then a subset  $I \subseteq V \setminus Z$  is independent in  $M/Z$  iff  $I \cup B_Z$  is independent in  $M$ .
- The rank function takes the form

$$r_{M/Z}(Y) = r(Y \cup Z) - r(Z) = r(Y|Z) \quad (8.13)$$

$$= r(Y \cup B_Z) - r(B_Z) = r(Y|B_Z) \quad (8.14)$$

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ . Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .
- Let  $Z \subseteq V$  and let  $B_Z$  be a base of  $Z$ . Then a subset  $I \subseteq V \setminus Z$  is independent in  $M/Z$  iff  $I \cup B_Z$  is independent in  $M$ .
- The rank function takes the form

$$r_{M/Z}(Y) = r(Y \cup Z) - r(Z) = r(Y|Z) \quad (8.13)$$

$$= r(Y \cup B_Z) - r(B_Z) = r(Y|B_Z) \quad (8.14)$$

- So given  $I \subseteq V \setminus Z$  and  $B_Z$  is a base of  $Z$ ,  $r_{M/Z}(I) = |I|$  is identical to  $r(I \cup Z) = |I| + r(Z) = |I| + |B_Z|$ . Since  $r(I \cup Z) = r(I \cup B_Z)$ , this implies  $r(I \cup B_Z) = |I| + |B_Z|$ , or  $I \cup B_Z$  is independent in  $M$ .

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ . Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .
- Let  $Z \subseteq V$  and let  $B_Z$  be a base of  $Z$ . Then a subset  $I \subseteq V \setminus Z$  is independent in  $M/Z$  iff  $I \cup B_Z$  is independent in  $M$ .
- The rank function takes the form

$$r_{M/Z}(Y) = r(Y \cup Z) - r(Z) = r(Y|Z) \quad (8.13)$$

$$= r(Y \cup B_Z) - r(B_Z) = r(Y|B_Z) \quad (8.14)$$

- So given  $I \subseteq V \setminus Z$  and  $B_Z$  is a base of  $Z$ ,  $r_{M/Z}(I) = |I|$  is identical to  $r(I \cup Z) = |I| + r(Z) = |I| + |B_Z|$ . Since  $r(I \cup Z) = r(I \cup B_Z)$ , this implies  $r(I \cup B_Z) = |I| + |B_Z|$ , or  $I \cup B_Z$  is independent in  $M$ .
- A **minor** of a matroid is any matroid obtained via a series of deletions and contractions of some matroid.

# Matroid contraction $M/Z$

- Contraction by  $Z$  is dual to deletion, and is like a forced inclusion of a contained base  $B_Z$  of  $Z$ , but with a similar ground set removal by  $Z$ . Contracting  $Z$  is written  $M/Z$ . Updated ground set in  $M/Z$  is  $V \setminus Z$ .
- Let  $Z \subseteq V$  and let  $B_Z$  be a base of  $Z$ . Then a subset  $I \subseteq V \setminus Z$  is independent in  $M/Z$  iff  $I \cup B_Z$  is independent in  $M$ .
- The rank function takes the form

$$r_{M/Z}(Y) = r(Y \cup Z) - r(Z) = r(Y|Z) \quad (8.13)$$

$$= r(Y \cup B_Z) - r(B_Z) = r(Y|B_Z) \quad (8.14)$$

- So given  $I \subseteq V \setminus Z$  and  $B_Z$  is a base of  $Z$ ,  $r_{M/Z}(I) = |I|$  is identical to  $r(I \cup Z) = |I| + r(Z) = |I| + |B_Z|$ . Since  $r(I \cup Z) = r(I \cup B_Z)$ , this implies  $r(I \cup B_Z) = |I| + |B_Z|$ , or  $I \cup B_Z$  is independent in  $M$ .
- A **minor** of a matroid is any matroid obtained via a series of deletions and contractions of some matroid.
- In fact, it is the case  $M/Z = (M^* \setminus Z)^*$  (Exercise: show why).



# Matroid Intersection

- Let  $M_1 = (V, \mathcal{I}_1)$  and  $M_2 = (V, \mathcal{I}_2)$  be two matroids. Consider their common independent sets  $\mathcal{I}_1 \cap \mathcal{I}_2$ .

# Matroid Intersection

- Let  $M_1 = (V, \mathcal{I}_1)$  and  $M_2 = (V, \mathcal{I}_2)$  be two matroids. Consider their common independent sets  $\mathcal{I}_1 \cap \mathcal{I}_2$ .
- While  $(V, \mathcal{I}_1 \cap \mathcal{I}_2)$  is typically not a matroid (**Exercise: show graphical example.**), we might be interested in finding the maximum size common independent set. That is, find  $\max |X|$  such that both  $X \in \mathcal{I}_1$  and  $X \in \mathcal{I}_2$ .

# Matroid Intersection

- Let  $M_1 = (V, \mathcal{I}_1)$  and  $M_2 = (V, \mathcal{I}_2)$  be two matroids. Consider their common independent sets  $\mathcal{I}_1 \cap \mathcal{I}_2$ .
- While  $(V, \mathcal{I}_1 \cap \mathcal{I}_2)$  is typically not a matroid (**Exercise: show graphical example.**), we might be interested in finding the maximum size common independent set. That is, find  $\max |X|$  such that both  $X \in \mathcal{I}_1$  and  $X \in \mathcal{I}_2$ .

## Theorem 8.4.1

Let  $M_1$  and  $M_2$  be given as above, with rank functions  $r_1$  and  $r_2$ . Then the size of the maximum size set in  $\mathcal{I}_1 \cap \mathcal{I}_2$  is given by

$$(r_1 * r_2)(V) \triangleq \min_{X \subseteq V} (r_1(X) + r_2(V \setminus X)) \quad (8.15)$$

# Matroid Intersection

- Let  $M_1 = (V, \mathcal{I}_1)$  and  $M_2 = (V, \mathcal{I}_2)$  be two matroids. Consider their common independent sets  $\mathcal{I}_1 \cap \mathcal{I}_2$ .
- While  $(V, \mathcal{I}_1 \cap \mathcal{I}_2)$  is typically not a matroid (**Exercise: show graphical example.**), we might be interested in finding the maximum size common independent set. That is, find  $\max |X|$  such that both  $X \in \mathcal{I}_1$  and  $X \in \mathcal{I}_2$ .

## Theorem 8.4.1

Let  $M_1$  and  $M_2$  be given as above, with rank functions  $r_1$  and  $r_2$ . Then the size of the maximum size set in  $\mathcal{I}_1 \cap \mathcal{I}_2$  is given by

$$(r_1 * r_2)(V) \triangleq \min_{X \subseteq V} (r_1(X) + r_2(V \setminus X)) \quad (8.15)$$

This is an instance of the **convolution of two submodular functions**,  $f_1$  and  $f_2$  that, evaluated at  $Y \subseteq V$ , is written as:

$$(f_1 * f_2)(Y) = \min_{X \subseteq Y} (f_1(X) + f_2(Y \setminus X)) \quad (8.16)$$

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$
- $\Leftrightarrow \min_X |\Gamma(X)| + |V| - |X| \geq |V|$



# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$
- $\Leftrightarrow \min_X |\Gamma(X)| + |V| - |X| \geq |V|$
- $\Leftrightarrow \min_X (|\Gamma(X)| + |V \setminus X|) \geq |V|$

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$
- $\Leftrightarrow \min_X |\Gamma(X)| + |V| - |X| \geq |V|$
- $\Leftrightarrow \min_X (|\Gamma(X)| + |V \setminus X|) \geq |V|$
- $\Leftrightarrow [\Gamma(\cdot) * |\cdot|](V) \geq |V|$

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$
- $\Leftrightarrow \min_X |\Gamma(X)| + |V| - |X| \geq |V|$
- $\Leftrightarrow \min_X (|\Gamma(X)| + |V \setminus X|) \geq |V|$
- $\Leftrightarrow [\Gamma(\cdot) * |\cdot|](V) \geq |V|$
- So Hall's theorem can be expressed as convolution. Exercise: define  $g(A) = [\Gamma(\cdot) * |\cdot|](A)$ , prove that  $g$  is submodular.

# Convolution and Hall's Theorem

- Recall Hall's theorem, that a transversal exists iff for all  $X \subseteq V$ , we have  $|\Gamma(X)| \geq |X|$ .
- $\Leftrightarrow |\Gamma(X)| - |X| \geq 0, \forall X$
- $\Leftrightarrow \min_X |\Gamma(X)| - |X| \geq 0$
- $\Leftrightarrow \min_X |\Gamma(X)| + |V| - |X| \geq |V|$
- $\Leftrightarrow \min_X (|\Gamma(X)| + |V \setminus X|) \geq |V|$
- $\Leftrightarrow [\Gamma(\cdot) * |\cdot|](V) \geq |V|$
- So Hall's theorem can be expressed as convolution. Exercise: define  $g(A) = [\Gamma(\cdot) * |\cdot|](A)$ , prove that  $g$  is submodular.
- Note, in general, convolution of two submodular functions does not preserve submodularity (but in certain special cases it does).

# Matroid Union

## Definition 8.4.2

Let  $M_1 = (V_1, \mathcal{I}_1)$ ,  $M_2 = (V_2, \mathcal{I}_2)$ ,  $\dots$ ,  $M_k = (V_k, \mathcal{I}_k)$  be matroids. We define the **union** of matroids as

$M_1 \vee M_2 \vee \dots \vee M_k = (V_1 \uplus V_2 \uplus \dots \uplus V_k, \mathcal{I}_1 \vee \mathcal{I}_2 \vee \dots \vee \mathcal{I}_k)$ , where

$$\mathcal{I}_1 \vee \mathcal{I}_2 \vee \dots \vee \mathcal{I}_k = \{I_1 \uplus I_2 \uplus \dots \uplus I_k \mid I_1 \in \mathcal{I}_1, \dots, I_k \in \mathcal{I}_k\} \quad (8.17)$$

Note  $A \uplus B$  designates the disjoint union of  $A$  and  $B$ .

# Matroid Union

## Definition 8.4.2

Let  $M_1 = (V_1, \mathcal{I}_1)$ ,  $M_2 = (V_2, \mathcal{I}_2)$ ,  $\dots$ ,  $M_k = (V_k, \mathcal{I}_k)$  be matroids. We define the **union** of matroids as

$M_1 \vee M_2 \vee \dots \vee M_k = (V_1 \uplus V_2 \uplus \dots \uplus V_k, \mathcal{I}_1 \vee \mathcal{I}_2 \vee \dots \vee \mathcal{I}_k)$ , where

$$\mathcal{I}_1 \vee \mathcal{I}_2 \vee \dots \vee \mathcal{I}_k = \{I_1 \uplus I_2 \uplus \dots \uplus I_k \mid I_1 \in \mathcal{I}_1, \dots, I_k \in \mathcal{I}_k\} \quad (8.17)$$

Note  $A \uplus B$  designates the disjoint union of  $A$  and  $B$ .

## Theorem 8.4.3

Let  $M_1 = (V_1, \mathcal{I}_1)$ ,  $M_2 = (V_2, \mathcal{I}_2)$ ,  $\dots$ ,  $M_k = (V_k, \mathcal{I}_k)$  be matroids, with rank functions  $r_1, \dots, r_k$ . Then the union of these matroids is still a matroid, having rank function

$$r(Y) = \min_{X \subseteq Y} \left( |Y \setminus X| + r_1(X \cap V_1) + \dots + r_k(X \cap V_k) \right) \quad (8.18)$$

for any  $Y \subseteq V_1 \uplus \dots \uplus V_2 \uplus \dots \uplus V_k$ .

# Exercise: Matroid Union, and Matroid duality

Exercise: Fully characterize  $M \vee M^*$ .

# Matroids of three or fewer elements are graphic

- All matroids up to and including three elements (edges) are graphic.



# Matroids of three or fewer elements are graphic


- All matroids up to and including three elements (edges) are graphic.




(a) The only matroid with zero elements.



(b) The two one-element matroids.






(c) The four two-element matroids.



(d) The eight three-element matroids.

# Matroids of three or fewer elements are graphic

- All matroids up to and including three elements (edges) are graphic.



(a) The only matroid with zero elements.

(b) The two one-element matroids.

(c) The four two-element matroids.

(d) The eight three-element matroids.

- This is a nice way to visualize matroids with very low ground set sizes. What about matroids that are low rank but with many elements?

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .
- Otherwise, the set is called **affinely independent**.

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .
- Otherwise, the set is called **affinely independent**.
- Concisely: points  $\{v_1, v_2, \dots, v_k\}$  are affinely independent if  $v_2 - v_1, v_3 - v_1, \dots, v_k - v_1$  are linearly independent.

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .
- Otherwise, the set is called **affinely independent**.
- Concisely: points  $\{v_1, v_2, \dots, v_k\}$  are affinely independent if  $v_2 - v_1, v_3 - v_1, \dots, v_k - v_1$  are linearly independent.
- Example: in 2D, three collinear points are affinely dependent, three non-collinear points are affinely independent, and  $\geq 4$  collinear or non-collinear points are affinely dependent.

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .
- Otherwise, the set is called **affinely independent**.
- Concisely: points  $\{v_1, v_2, \dots, v_k\}$  are affinely independent if  $v_2 - v_1, v_3 - v_1, \dots, v_k - v_1$  are linearly independent.
- Example: in 2D, three collinear points are affinely dependent, three non-collinear points are affinely independent, and  $\geq 4$  collinear or non-collinear points are affinely dependent.

## Proposition 8.5.1 (affine matroid)

Let ground set  $E = \{1, \dots, m\}$  index column vectors of a matrix, and let  $\mathcal{I}$  be the set of subsets  $X$  of  $E$  such that  $X$  indices affinely independent vectors. Then  $(E, \mathcal{I})$  is a matroid.

# Affine Matroids

- Given an  $n \times m$  matrix with entries over some field  $\mathbb{F}$ , we say that a subset  $S \subseteq \{1, \dots, m\}$  of indices (with corresponding column vectors  $\{v_i : i \in S\}$ , with  $|S| = k \leq m$ ) is **affinely dependent** if  $m \geq 1$  and there exists elements  $\{a_1, \dots, a_k\} \in \mathbb{F}$ , not all zero with  $\sum_{i=1}^k a_i = 0$ , such that  $\sum_{i=1}^k a_i v_i = 0$ .
- Otherwise, the set is called **affinely independent**.
- Concisely: points  $\{v_1, v_2, \dots, v_k\}$  are affinely independent if  $v_2 - v_1, v_3 - v_1, \dots, v_k - v_1$  are linearly independent.
- Example: in 2D, three collinear points are affinely dependent, three non-collinear points are affinely independent, and  $\geq 4$  collinear or non-collinear points are affinely dependent.

## Proposition 8.5.1 (affine matroid)

Let ground set  $E = \{1, \dots, m\}$  index column vectors of a matrix, and let  $\mathcal{I}$  be the set of subsets  $X$  of  $E$  such that  $X$  indices affinely independent vectors. Then  $(E, \mathcal{I})$  is a matroid.

**Exercise: prove this.**




# Euclidean Representation of Low-rank Matroids

- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .


# Euclidean Representation of Low-rank Matroids

- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .
- We can plot the points in  $\mathbb{R}^2$  as on the right:



# Euclidean Representation of Low-rank Matroids


- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .
- We can plot the points in  $\mathbb{R}^2$  as on the right:
- A point has rank 1, points that comprise a line have rank 2, points that comprise a plane have rank 3.






# Euclidean Representation of Low-rank Matroids

- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .
- We can plot the points in  $\mathbb{R}^2$  as on the right:
- A point has rank 1, points that comprise a line have rank 2, points that comprise a plane have rank 3.
- Flats (points, lines, planes, etc.) have rank equal to one more than their geometric dimension.
- Any two distinct points constitute a line, but lines with only two points are not drawn.




# Euclidean Representation of Low-rank Matroids

- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .
- We can plot the points in  $\mathbb{R}^2$  as on the right:
- A point has rank 1, points that comprise a line have rank 2, points that comprise a plane have rank 3.
- Flats (points, lines, planes, etc.) have rank equal to one more than their geometric dimension.
- Any two distinct points constitute a line, but lines with only two points are not drawn.
- Lines indicate collinear sets with  $\geq 3$  points, while any two points have rank 2.




# Euclidean Representation of Low-rank Matroids

- Consider the affine matroid with  $n \times m = 2 \times 6$  matrix on the field  $\mathbb{F} = \mathbb{R}$ , and let the elements be  $\{(0, 0), (1, 0), (2, 0), (0, 1), (0, 2), (1, 1)\}$ .
- We can plot the points in  $\mathbb{R}^2$  as on the right:
- A point has rank 1, points that comprise a line have rank 2, points that comprise a plane have rank 3.
- Flats (points, lines, planes, etc.) have rank equal to one more than their geometric dimension.
- Any two distinct points constitute a line, but lines with only two points are not drawn.
- Lines indicate collinear sets with  $\geq 3$  points, while any two points have rank 2.
- Dependent sets consist of all subsets with  $\geq 4$  elements (rank 3), or 3 collinear elements (rank 2).  
Any two points have rank 2.



# Euclidean Representation of Low-rank Matroids


- As another example on the right, a rank 4 matroid





# Euclidean Representation of Low-rank Matroids

- As another example on the right, a rank 4 matroid



- All sets of 5 points are dependent. The only other sets of dependent points are coplanar ones of size 4. Namely:
  - $\{(0, 0, 0), (0, 1, 0), (1, 1, 0), (1, 0, 0)\}$ ,
  - $\{(0, 0, 0), (0, 0, 1), (0, 1, 1), (0, 1, 0)\}$ , and
  - $\{(0, 0, 1), (0, 1, 1), (1, 1, 0), (1, 0, 0)\}$ .

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - ①  $|X| \geq 2$  and the points are identical;

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - 1  $|X| \geq 2$  and the points are identical;
  - 2  $|X| \geq 3$  and the points are collinear;

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - 1  $|X| \geq 2$  and the points are identical;
  - 2  $|X| \geq 3$  and the points are collinear;
  - 3  $|X| \geq 4$  and the points are coplanar; or

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - 1  $|X| \geq 2$  and the points are identical;
  - 2  $|X| \geq 3$  and the points are collinear;
  - 3  $|X| \geq 4$  and the points are coplanar; or
  - 4  $|X| \geq 5$  and the points are anywhere in space.

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - ①  $|X| \geq 2$  and the points are identical;
  - ②  $|X| \geq 3$  and the points are collinear;
  - ③  $|X| \geq 4$  and the points are coplanar; or
  - ④  $|X| \geq 5$  and the points are anywhere in space.
- When they exist, loops are represented in a geometry by a separate box indicating how many loops there are.

# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - ①  $|X| \geq 2$  and the points are identical;
  - ②  $|X| \geq 3$  and the points are collinear;
  - ③  $|X| \geq 4$  and the points are coplanar; or
  - ④  $|X| \geq 5$  and the points are anywhere in space.
- When they exist, loops are represented in a geometry by a separate box indicating how many loops there are.
- Parallel elements, when they exist in a matroid, are indicated by a multiplicity next to a point.

True regardless of how big  $|V|$  is.



# Euclidean Representation of Low-rank Matroids

- In general, for a matroid  $\mathcal{M}$  of rank  $m + 1$  with  $m \leq 3$ , then a subset  $X$  in a geometric representation in  $\mathbb{R}^m$  is dependent if:
  - ①  $|X| \geq 2$  and the points are identical;
  - ②  $|X| \geq 3$  and the points are collinear;
  - ③  $|X| \geq 4$  and the points are coplanar; or
  - ④  $|X| \geq 5$  and the points are anywhere in space.
- When they exist, loops are represented in a geometry by a separate box indicating how many loops there are.
- Parallel elements, when they exist in a matroid, are indicated by a multiplicity next to a point.

## Theorem 8.5.2

*Any matroid of rank  $m \leq 4$  can be represented by an affine matroid in  $\mathbb{R}^{m-1}$ .*

True regardless of how big  $|V|$  is.

# Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)
- every plane contains at least three non-collinear points (not dependent unless  $> 3$ )

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)
- every plane contains at least three non-collinear points (not dependent unless  $> 3$ )
- any three distinct non-collinear points lie on a plane

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)
- every plane contains at least three non-collinear points (not dependent unless  $> 3$ )
- any three distinct non-collinear points lie on a plane
- If diagram has at most one plane, then any two distinct lines meet in at most one point.



# Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)
- every plane contains at least three non-collinear points (not dependent unless  $> 3$ )
- any three distinct non-collinear points lie on a plane
- If diagram has at most one plane, then any two distinct lines meet in at most one point.
- If diagram has more than one plane, then: 1) any two distinct planes meeting in more than two points do so in a line; 2) any two distinct lines meeting in a point do so in at most one point and lie in on a common plane; 3) any line not lying on a plane intersects it in at most one point.

## Euclidean Rep. of Low-rank Matroids: Conditions

- rank-1 (resp. rank-2, rank-3) flats correspond to points (resp. lines, planes).
- a set of parallel points (could be size 1) does not touch another set of parallel points (could be size 1).
- every line contains at least two points (not dependent unless  $> 2$ ).
- any two distinct points lie on a line (often not drawn when only two)
- every plane contains at least three non-collinear points (not dependent unless  $> 3$ )
- any three distinct non-collinear points lie on a plane
- If diagram has at most one plane, then any two distinct lines meet in at most one point.
- If diagram has more than one plane, then: 1) any two distinct planes meeting in more than two points do so in a line; 2) any two distinct lines meeting in a point do so in at most one point and lie in on a common plane; 3) any line not lying on a plane intersects it in at most one point.
- (see Oxley 2011 for more details).

# Euclidean Representation of Low-rank Matroids


- Very useful for graphically depicting low-rank matrices but which still have rich structure. Also useful for answering questions.

# Euclidean Representation of Low-rank Matroids

- Very useful for graphically depicting low-rank matrices but which still have rich structure. Also useful for answering questions.
- Example: Is there a matroid that is not representable (i.e., not linear for some field)?


# Euclidean Representation of Low-rank Matroids

- Very useful for graphically depicting low-rank matrices but which still have rich structure. Also useful for answering questions.
- Example: Is there a matroid that is not representable (i.e., not linear for some field)? **Yes, consider the matroid**



# Euclidean Representation of Low-rank Matroids


- Very useful for graphically depicting low-rank matrices but which still have rich structure. Also useful for answering questions.
- Example: Is there a matroid that is not representable (i.e., not linear for some field)? Yes, consider the matroid



- Called the non-Pappus matroid. Has rank three, but any matrix matroid with the above dependencies would require that  $\{7, 8, 9\}$  is dependent, hence requiring an additional line in the above.


# Euclidean Representation of Low-rank Matroids: A test

- Is this a matroid?



# Euclidean Representation of Low-rank Matroids: A test

- Is this a matroid?




- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ .  
So  $r(X) =$



# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ .  
So  $r(X) = 3$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ .  
So  $r(X) = 3$ , and  $r(Y) =$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ .  
So  $r(X) = 3$ , and  $r(Y) = 3$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) =$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) = 4$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) = 4$ , so we must have, by submodularity, that
 
$$r(\{1, 6, 7\}) = r(X \cap Y) \leq r(X) + r(Y) - r(X \cup Y) = 2.$$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) = 4$ , so we must have, by submodularity, that
 
$$r(\{1, 6, 7\}) = r(X \cap Y) \leq r(X) + r(Y) - r(X \cup Y) = 2.$$
- However, from the diagram, we have that since 1, 6, 7 are distinct non-collinear points, we have that  $r(X \cap Y) =$

# Euclidean Representation of Low-rank Matroids: A test

- Is this a matroid?




- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) = 4$ , so we must have, by submodularity, that
 
$$r(\{1, 6, 7\}) = r(X \cap Y) \leq r(X) + r(Y) - r(X \cup Y) = 2.$$
- However, from the diagram, we have that since 1, 6, 7 are distinct non-collinear points, we have that  $r(X \cap Y) = 3$



# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- Check rank's submodularity: Let  $X = \{1, 2, 3, 6, 7\}$ ,  $Y = \{1, 4, 5, 6, 7\}$ . So  $r(X) = 3$ , and  $r(Y) = 3$ , and  $r(X \cup Y) = 4$ , so we must have, by submodularity, that
 
$$r(\{1, 6, 7\}) = r(X \cap Y) \leq r(X) + r(Y) - r(X \cup Y) = 2.$$
- However, from the diagram, we have that since 1, 6, 7 are distinct non-collinear points, we have that  $r(X \cap Y) = 3$

# Euclidean Representation of Low-rank Matroids: A test


- Is this a matroid?



- If we extend the line from 6-7 to 1, then is it a matroid?
- Hence, not all 2D or 3D graphs of points and lines are matroids.


# Matroid?

- Consider the following geometry on  $|V| = 8$  points with  $V = \{a, b, c, d, e, f, g, h\}$ .



# Matroid?


- Consider the following geometry on  $|V| = 8$  points with  $V = \{a, b, c, d, e, f, g, h\}$ .



- Note, we are given that the points  $\{b, d, h, f\}$  are not coplanar. However, the following sets of points are coplanar:  $\{a, b, e, f\}$ ,  $\{d, c, g, h\}$ ,  $\{a, d, h, e\}$ ,  $\{b, c, g, f\}$ ,  $\{b, c, d, a\}$ ,  $\{f, g, h, e\}$ , and  $\{a, c, g, e\}$ .

# Matroid?



- Consider the following geometry on  $|V| = 8$  points with  $V = \{a, b, c, d, e, f, g, h\}$ .



- Note, we are given that the points  $\{b, d, h, f\}$  are not coplanar. However, the following sets of points are coplanar:  $\{a, b, e, f\}$ ,  $\{d, c, g, h\}$ ,  $\{a, d, h, e\}$ ,  $\{b, c, g, f\}$ ,  $\{b, c, d, a\}$ ,  $\{f, g, h, e\}$ , and  $\{a, c, g, e\}$ .
- Exercise: Is this a matroid? Exercise: If so, is it representable?**

# Projective Geometries: Other Examples



- Other examples can be more complex, consider the following two matroids (from Oxley, 2011):





# Projective Geometries: Other Examples

- Other examples can be more complex, consider the following two matroids (from Oxley, 2011):



- Right: a matroid (and a 2D depiction of a geometry) over the field  $GF(3) = \{0, 1, 2\} \pmod{3}$  and is “coordinatizable” in  $GF(3)^3$ .
- Hence, lines (in 2D) which are rank 2 sets may be curved; planes (in 3D) can be twisted.



# Matroids, Representation and Equivalence: Summary

- Matroids with  $|V| \leq 3$  are graphic.

# Matroids, Representation and Equivalence: Summary

- Matroids with  $|V| \leq 3$  are graphic.
- Matroids with  $r(V) \leq 4$  can be geometrically represented in  $\mathbb{R}^3$ .

# Matroids, Representation and Equivalence: Summary

- Matroids with  $|V| \leq 3$  are graphic.
- Matroids with  $r(V) \leq 4$  can be geometrically represented in  $\mathbb{R}^3$ .
- Not all matroids are linear (i.e., matric) matroids.

# Matroids, Representation and Equivalence: Summary

- Matroids with  $|V| \leq 3$  are graphic.
- Matroids with  $r(V) \leq 4$  can be geometrically represented in  $\mathbb{R}^3$ .
- Not all matroids are linear (i.e., matric) matroids.
- Matroids can be seen as related to projective geometries (and are sometimes called combinatorial geometries).

# Matroids, Representation and Equivalence: Summary

- Matroids with  $|V| \leq 3$  are graphic.
- Matroids with  $r(V) \leq 4$  can be geometrically represented in  $\mathbb{R}^3$ .
- Not all matroids are linear (i.e., matric) matroids.
- Matroids can be seen as related to projective geometries (and are sometimes called combinatorial geometries).
- Exists much research on different subclasses of matroids, and if/when they are contained in (or isomorphic to) each other.

# Matroid Further Reading

- “Matroids: A Geometric Introduction”, Gordon and McNulty, 2012.
- “The Coming of the Matroids”, William Cunningham, 2012 (a nice history)
- Welsh, “Matroid Theory”, 1975.
- Oxley, “Matroid Theory”, 1992 (and 2011) (perhaps best “single source” on matroids right now).
- Crapo & Rota, “On the Foundations of Combinatorial Theory: Combinatorial Geometries”, 1970 (while this is old, it is very readable).
- Lawler, “Combinatorial Optimization: Networks and Matroids”, 1976.
- Schrijver, “Combinatorial Optimization”, 2003

# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.

# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.
- The goal is to choose a good subset of items, and the fundamental tenet of the greedy algorithm is to **choose next whatever currently looks best**, without the possibility of later recall or backtracking.



# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.
- The goal is to choose a good subset of items, and the fundamental tenet of the greedy algorithm is to **choose next whatever currently looks best**, without the possibility of later recall or backtracking.
- Sometimes, this gives the optimal solution (we saw three greedy algorithms that can find the maximum weight spanning tree).

# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.
- The goal is to choose a good subset of items, and the fundamental tenet of the greedy algorithm is to **choose next whatever currently looks best**, without the possibility of later recall or backtracking.
- Sometimes, this gives the optimal solution (we saw three greedy algorithms that can find the maximum weight spanning tree).
- Greedy is good since it can be made to run very fast  $O(n \log n)$ .

# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.
- The goal is to choose a good subset of items, and the fundamental tenet of the greedy algorithm is to **choose next whatever currently looks best**, without the possibility of later recall or backtracking.
- Sometimes, this gives the optimal solution (we saw three greedy algorithms that can find the maximum weight spanning tree).
- Greedy is good since it can be made to run very fast  $O(n \log n)$ .
- Often, however, greedy is heuristic (it might work well in practice, but worst-case performance can be unboundedly poor).

# The greedy algorithm

- In combinatorial optimization, the greedy algorithm is often useful as a heuristic that can work quite well in practice.
- The goal is to choose a good subset of items, and the fundamental tenet of the greedy algorithm is to **choose next whatever currently looks best**, without the possibility of later recall or backtracking.
- Sometimes, this gives the optimal solution (we saw three greedy algorithms that can find the maximum weight spanning tree).
- Greedy is good since it can be made to run very fast  $O(n \log n)$ .
- Often, however, greedy is heuristic (it might work well in practice, but worst-case performance can be unboundedly poor).
- We will next see that the greedy algorithm working optimally is a defining property of a matroid, and is also a defining property of a polymatroid function.

# Matroid and the greedy algorithm

- Let  $(E, \mathcal{I})$  be an independence system, and we are given a non-negative modular weight function  $w : E \rightarrow \mathbb{R}_+$ .

# Matroid and the greedy algorithm

- Let  $(E, \mathcal{I})$  be an independence system, and we are given a non-negative modular weight function  $w : E \rightarrow \mathbb{R}_+$ .

---

## Algorithm 1: The Matroid Greedy Algorithm

---

- 1 Set  $X \leftarrow \emptyset$  ;
  - 2 **while**  $\exists v \in E \setminus X$  s.t.  $X \cup \{v\} \in \mathcal{I}$  **do**
  - 3      $v \in \operatorname{argmax} \{w(v) : v \in E \setminus X, X \cup \{v\} \in \mathcal{I}\}$  ;
  - 4      $X \leftarrow X \cup \{v\}$  ;
-

# Matroid and the greedy algorithm

- Let  $(E, \mathcal{I})$  be an independence system, and we are given a non-negative modular weight function  $w : E \rightarrow \mathbb{R}_+$ .

---

## Algorithm 1: The Matroid Greedy Algorithm

---

- 1 Set  $X \leftarrow \emptyset$  ;
  - 2 **while**  $\exists v \in E \setminus X$  s.t.  $X \cup \{v\} \in \mathcal{I}$  **do**
  - 3      $v \in \operatorname{argmax} \{w(v) : v \in E \setminus X, X \cup \{v\} \in \mathcal{I}\}$  ;
  - 4      $X \leftarrow X \cup \{v\}$  ;
- 
- Same as sorting items by decreasing weight  $w$ , and then choosing items in that order that retain independence.

# Matroid and the greedy algorithm

- Let  $(E, \mathcal{I})$  be an independence system, and we are given a non-negative modular weight function  $w : E \rightarrow \mathbb{R}_+$ .

---

## Algorithm 1: The Matroid Greedy Algorithm

---

- 1 Set  $X \leftarrow \emptyset$  ;
  - 2 **while**  $\exists v \in E \setminus X$  s.t.  $X \cup \{v\} \in \mathcal{I}$  **do**
  - 3      $v \in \operatorname{argmax} \{w(v) : v \in E \setminus X, X \cup \{v\} \in \mathcal{I}\}$  ;
  - 4      $X \leftarrow X \cup \{v\}$  ;
- 
- Same as sorting items by decreasing weight  $w$ , and then choosing items in that order that retain independence.

### Theorem 8.6.1

Let  $(E, \mathcal{I})$  be an independence system. Then the pair  $(E, \mathcal{I})$  is a matroid *if and only if* for each weight function  $w \in \mathcal{R}_+^E$ , Algorithm 1 above leads to a set  $I \in \mathcal{I}$  of maximum weight  $w(I)$ .



# Review from Lecture 6

- The next slide is from Lecture 6.

# Matroids by bases

In general, besides independent sets and rank functions, there are other equivalent ways to characterize matroids.

## Theorem 8.6.3 (Matroid (by bases))

Let  $E$  be a set and  $\mathcal{B}$  be a nonempty collection of subsets of  $E$ . Then the following are equivalent.

- ①  $\mathcal{B}$  is the collection of bases of a matroid;
- ② if  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B' - x + y \in \mathcal{B}$  for some  $y \in B \setminus B'$ .
- ③ If  $B, B' \in \mathcal{B}$ , and  $x \in B' \setminus B$ , then  $B - y + x \in \mathcal{B}$  for some  $y \in B \setminus B'$ .

Properties 2 and 3 are called “exchange properties.”

Proof here is omitted but think about this for a moment in terms of linear spaces and matrices, and (alternatively) spanning trees.

# Matroid and the greedy algorithm

proof of Theorem 8.6.1.

- Assume  $(E, \mathcal{I})$  is a matroid and  $w : E \rightarrow \mathcal{R}_+$  is given.

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume  $(E, \mathcal{I})$  is a matroid and  $w : E \rightarrow \mathcal{R}_+$  is given.
- Let  $A = (a_1, a_2, \dots, a_r)$  be the solution returned by greedy, where  $r = r(M)$  the rank of the matroid, and we order the elements as they were chosen (so  $w(a_1) \geq w(a_2) \geq \dots \geq w(a_r)$ ).

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume  $(E, \mathcal{I})$  is a matroid and  $w : E \rightarrow \mathcal{R}_+$  is given.
- Let  $A = (a_1, a_2, \dots, a_r)$  be the solution returned by greedy, where  $r = r(M)$  the rank of the matroid, and we order the elements as they were chosen (so  $w(a_1) \geq w(a_2) \geq \dots \geq w(a_r)$ ).
- $A$  is a base of  $M$ , and let  $B = (b_1, \dots, b_r)$  be any another base of  $M$  with elements also ordered decreasing by weight, so  $w(b_1) \geq w(b_2) \geq \dots \geq w(b_r)$ .

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume  $(E, \mathcal{I})$  is a matroid and  $w : E \rightarrow \mathcal{R}_+$  is given.
- Let  $A = (a_1, a_2, \dots, a_r)$  be the solution returned by greedy, where  $r = r(M)$  the rank of the matroid, and we order the elements as they were chosen (so  $w(a_1) \geq w(a_2) \geq \dots \geq w(a_r)$ ).
- $A$  is a base of  $M$ , and let  $B = (b_1, \dots, b_r)$  be any another base of  $M$  with elements also ordered decreasing by weight, so  $w(b_1) \geq w(b_2) \geq \dots \geq w(b_r)$ .
- We next show that not only is  $w(A) \geq w(B)$  but that  $w(a_i) \geq w(b_i)$  for all  $i$ .

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume otherwise, and let  $k$  be the first (smallest) integer such that  $w(a_k) < w(b_k)$ . Hence  $w(a_j) \geq w(b_j)$  for  $j < k$ .

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume otherwise, and let  $k$  be the first (smallest) integer such that  $w(a_k) < w(b_k)$ . Hence  $w(a_j) \geq w(b_j)$  for  $j < k$ .
- Define independent sets  $A_{k-1} = \{a_1, \dots, a_{k-1}\}$  and  $B_k = \{b_1, \dots, b_k\}$ .

...



# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume otherwise, and let  $k$  be the first (smallest) integer such that  $w(a_k) < w(b_k)$ . Hence  $w(a_j) \geq w(b_j)$  for  $j < k$ .
- Define independent sets  $A_{k-1} = \{a_1, \dots, a_{k-1}\}$  and  $B_k = \{b_1, \dots, b_k\}$ .
- Since  $|A_{k-1}| < |B_k|$ , there exists a  $b_i \in B_k \setminus A_{k-1}$  where  $A_{k-1} \cup \{b_i\} \in \mathcal{I}$  for some  $1 \leq i \leq k$ .

...

# Matroid and the greedy algorithm

## proof of Theorem 8.6.1.

- Assume otherwise, and let  $k$  be the first (smallest) integer such that  $w(a_k) < w(b_k)$ . Hence  $w(a_j) \geq w(b_j)$  for  $j < k$ .
- Define independent sets  $A_{k-1} = \{a_1, \dots, a_{k-1}\}$  and  $B_k = \{b_1, \dots, b_k\}$ .
- Since  $|A_{k-1}| < |B_k|$ , there exists a  $b_i \in B_k \setminus A_{k-1}$  where  $A_{k-1} \cup \{b_i\} \in \mathcal{I}$  for some  $1 \leq i \leq k$ .
- But  $w(b_i) \geq w(b_k) > w(a_k)$ , and so the greedy algorithm would have chosen  $b_i$  rather than  $a_k$ , contradicting what greedy does.



# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Given an independence system  $(E, \mathcal{I})$ , suppose the greedy algorithm leads to an independent set of max weight for every non-negative weight function. We'll show  $(E, \mathcal{I})$  is a matroid.

# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Given an independence system  $(E, \mathcal{I})$ , suppose the greedy algorithm leads to an independent set of max weight for every non-negative weight function. We'll show  $(E, \mathcal{I})$  is a matroid.
- Emptyset containing and down monotonicity already holds (since we've started with an independence system).

# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Given an independence system  $(E, \mathcal{I})$ , suppose the greedy algorithm leads to an independent set of max weight for every non-negative weight function. We'll show  $(E, \mathcal{I})$  is a matroid.
- Emptyset containing and down monotonicity already holds (since we've started with an independence system).
- Let  $I, J \in \mathcal{I}$  with  $|I| < |J|$ . Suppose to the contrary, that  $I \cup \{z\} \notin \mathcal{I}$  for all  $z \in J \setminus I$ .

# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Given an independence system  $(E, \mathcal{I})$ , suppose the greedy algorithm leads to an independent set of max weight for every non-negative weight function. We'll show  $(E, \mathcal{I})$  is a matroid.
- Emptyset containing and down monotonicity already holds (since we've started with an independence system).
- Let  $I, J \in \mathcal{I}$  with  $|I| < |J|$ . Suppose to the contrary, that  $I \cup \{z\} \notin \mathcal{I}$  for all  $z \in J \setminus I$ .
- Define the following modular weight function  $w$  on  $E$ , and define  $k = |I|$ .

$$w(v) = \begin{cases} k + 2 & \text{if } v \in I, \\ k + 1 & \text{if } v \in J \setminus I, \\ 0 & \text{if } v \in E \setminus (I \cup J) \end{cases} \quad (8.19)$$

# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Now greedy will, after  $k$  iterations, recover  $I$ , but it cannot choose any element in  $J \setminus I$  by assumption. Thus, greedy chooses a set of weight  $k(k+2) = w(I)$ .

# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Now greedy will, after  $k$  iterations, recover  $I$ , but it cannot choose any element in  $J \setminus I$  by assumption. Thus, greedy chooses a set of weight  $k(k+2) = w(I)$ .
- On the other hand,  $J$  has weight

$$w(J) \geq |J|(k+1) \geq (k+1)(k+1) > k(k+2) = w(I) \quad (8.20)$$

so  $J$  has strictly larger weight but is still independent, contradicting greedy's optimality.



# Matroid and the greedy algorithm

## converse proof of Theorem 8.6.1.

- Now greedy will, after  $k$  iterations, recover  $I$ , but it cannot choose any element in  $J \setminus I$  by assumption. Thus, greedy chooses a set of weight  $k(k+2) = w(I)$ .
- On the other hand,  $J$  has weight

$$w(J) \geq |J|(k+1) \geq (k+1)(k+1) > k(k+2) = w(I) \quad (8.20)$$

so  $J$  has strictly larger weight but is still independent, contradicting greedy's optimality.

- Therefore, there must be a  $z \in J \setminus I$  such that  $I \cup \{z\} \in \mathcal{I}$ , and since  $I$  and  $J$  are arbitrary,  $(E, \mathcal{I})$  must be a matroid.

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.
- If we don't want elements with weight 0, we can stop once (and if) the weight hits zero, thus giving us a maximum weight independent set.

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.
- If we don't want elements with weight 0, we can stop once (and if) the weight hits zero, thus giving us a maximum weight independent set.
- We don't need non-negativity, we can use any  $w \in \mathbb{R}^E$  and keep going until we have a base.

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.
- If we don't want elements with weight 0, we can stop once (and if) the weight hits zero, thus giving us a maximum weight independent set.
- We don't need non-negativity, we can use any  $w \in \mathbb{R}^E$  and keep going until we have a base.
- If we stop at a negative value, we'll once again get a maximum weight independent set.

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.
- If we don't want elements with weight 0, we can stop once (and if) the weight hits zero, thus giving us a maximum weight independent set.
- We don't need non-negativity, we can use any  $w \in \mathbb{R}^E$  and keep going until we have a base.
- If we stop at a negative value, we'll once again get a maximum weight independent set.
- **Exercise: what if we keep going until a base even if we encounter negative values?**

# Matroid and greedy

- As given, the theorem asked for a modular function  $w \in \mathbb{R}_+^E$ .
- This will not only return an independent set, but it will return a base if we keep going even if the weights are 0.
- If we don't want elements with weight 0, we can stop once (and if) the weight hits zero, thus giving us a maximum weight independent set.
- We don't need non-negativity, we can use any  $w \in \mathbb{R}^E$  and keep going until we have a base.
- If we stop at a negative value, we'll once again get a maximum weight independent set.
- **Exercise: what if we keep going until a base even if we encounter negative values?**
- We can instead do **as small as possible** thus giving us a minimum weight independent set/base.



# Summary of Important (for us) Matroid Definitions

Given an independence system, matroids are defined equivalently by any of the following:

- All maximally independent sets have the same size.
- A monotone non-decreasing submodular integral rank function with unit increments.
- The greedy algorithm achieves the maximum weight independent set for all weight functions.